

**ALGORITHMIC DETERMINATION OF FLANKING DNA SEQUENCES
THAT CONTROL THE EXPRESSION OF SETS OF GENES IN
PROKARYOTIC, ARCHEA AND EUKARYOTIC GENOMES**

5 Reference to Related Application

The present application is the subject of Provisional Application Serial No. 60/208,650 filed June 2, 2000 entitled ALGORITHMIC DETERMINATION OF CONNECTRONS FOR THE HIGH LEVEL REGULATION OF GENE EXPRESSION.

10 Introduction

RNA introduced into a cell by a virus is now known to trigger a cellular defense mechanism known as post-transcriptional gene silencing (PTGS). If the viral RNA sequence matches a sequence within the cell's genome the associated genes are turned off or silenced. This phenomenon is also called 'RNA interference' or RNAi. A single-stranded RNA can interact with another single-stranded RNA (known as antisense RNA). The single-stranded RNA can also form a triple-stranded complex with double-stranded DNA. This triple-stranded complex is known as a Hoogsteen helix. This patent application shows how two specific adjacent RNA single-stranded sequences (called C1 and C2 – for Control Sequence 1 and Control Sequence 2) interact with two distant double-stranded DNA sequences (called T1 and T2 – for Target Sequence 1 and Target Sequence 2) to form a tetradic relationship which is called a "connectron". The two distant DNA double-stranded sequences (T1 and T2) must be on the same chromosome in a genome and they must be between about 1kb and 105kb of each other. The adjacent single-stranded RNA sequences (C1/C2) can be on the same or different chromosome as the T1 and T2 sequences. The C1 sequence is identical to the T1 sequence and the C2 sequence is identical to the T2

sequence. The connectron acts to stabilize the double-stranded DNA by allowing 30nm chromatin particles to form. Genes that lie between the T1 and T2 sequences when wrapped up in 30nm chromatin particles are not open to promotion and expression. The connectron (i.e. the tetradic relationship between the T1-T2 sequences and C1/C2 sequences) provides a general explanation for PTGS. A connectron can be implemented by RNA sequences, PNA (Peptide Nucleic Acid) sequences or by a zinc-finger DNA Binding Protein (DBP) specific to the T1 and T2 sequences.

Characteristically the adjacent C1/C2 sequences lie in the 3'UTR of a gene. The T1 and T2 sequences do not lie within the translated region of any gene. These sequences "surround" one or more genes. There are, however, T1 and T2 sequence pairs that surround one or more C1/C2 sequences that are not 3'UTR to any gene. These are called "geneless connectrons". There may be promoter sequences that cause the transcription of these 3'UTR sequences.

A computer-based algorithm that is similar to the algorithm used in the US Patent 6,205,404 has been developed to determine the connectron structure of any genome. This algorithm determines the existence of all the connectrons in the genomic DNA. Connectrons exist in prokaryotes, archaea, single-celled eukaryotes, multi-celled eukaryotes, plants and higher animals. Connectron relationships exist between prokaryotes and their plasmids. The geneless connectrons provide a possible mechanism for forming a hierarchy of gene expression control that will produce an understanding of cell differentiation and tissue development.

Each connectron is a unique tetrad of sequences. Each connectron changes the expression of the genes between the T1 and T2 sequences. The C1 sequence (which is equivalent to the T1 sequence) and the C2 sequence (which is equivalent to the T2 sequence) are determined by the invention described in this patent application. In general, the tetrad of connectron sequences can be patented because the structure of matter is known and the function of specific gene expression modulation is also known. Gene expression modification can be produced by introducing antisense

RNA or PNA to interact C1/C2 RNA sequences or zinc-finger DBPs to interact with the T1 and T2 sequences. Using connectrons it will be possible to modify cellular and tissue behavior in a very general manner.

Examples will be given from different genomes to illustrate that the connectron is a perfectly general and universal concept.

Definitions

Double stranded DNA – Watson and Crick showed in 1953 that DNA naturally forms a double-stranded helix. A typical double stranded sequence is

5'-TAGAGGAGTACCAC-3'

3'-ATCTCCTCATGGTG-5'

Hydrogen Bond - The force between a hydrogen atom and another heavier atom such as Oxygen (O), Nitrogen (N), Phosphorus (P), or Sulfur (S).

Positive strand – The positive strand is normally represented 5' to 3' running left to right as in

5'-TAGAGGAGTACCAC-3'

Negative strand – The negative strand is normally represented 5' to 3' running right to left as in

3'-ATCTCCTCATGGTG-5'

Single stranded RNA – Either the positive or the negative strand of the double-stranded DNA can be transcribed by the polymerase. In RNA U replaces T.

5 RNA of positive strand sequence 5'-UAGAGGAGUACCAC-3'
RNA of negative strand sequence 5'-GUGGUACUCCUCUA-3'

Antisense RNA – The antisense strand of any RNA sequence is the complement sequence

10

RNA sequence 5'-UAGAGGAGUACCAC-3'
Antisense RNA sequence 3'-AUCUCCUCAUGGUG-5'

Triple Strand Helix – The RNA sequence of a RNA/DNA triple-strand complex is the same as the positive strand of the DNA

15

DNA positive strand 5'-TAGAGGAGTACCAC-3'
DNA negative strand 3'-ATCTCCTCATGGTG-5'
RNA strand 5'-UAGAGGAGUACCAC-3'

20

Promoter – Any region of DNA, that binds proteins which engage the polymerase transcription mechanism.

TATA Box – A region near the 3' end of a promoter with the sequence TATA.

25

mRNA – The RNA produced from the DNA by the polymerase as a result of transcription

Start of transcription – The 3' end of a promoter where the polymerase mechanism begins to transcribe DNA into mRNA.

30

Exon – Any region of mRNA which is used to code for proteins

Intron – Any region of mRNA lying between two exons which is not used to code for proteins. The introns are edited out of the initial RNA transcript to form the mature mRNA.

5

3' UTR – The untranslated 3' end of an mRNA is beyond the end of the last exon. A stop codon in the mRNA causes the ribosome to stop the translation of mRNA into protein.


10

End of translation – The 3' end of the 3'-most exon.

Translated region – Any collection of exons and introns.

Gene – Any DNA region that codes for a protein. Introns do not occur in prokaryotic genes and they sometime fail to occur in eukaryotic genes. A typical model of a gene is

15



20

25

Positive strand gene – Any gene in which the features run 5' to 3' on the positive strand

30

Negative strand gene – Any gene in which the features run 5' to 3' on the negative strand

35

C1 sequence – Any positive or negative strand DNA sequence of 20 bases or more.

The C2 sequence must occur in the same chromosome as the C1 sequence.

C2 sequence – Any positive or negative strand DNA sequence of 20 bases or more.

The C1 sequence must occur in the same chromosome as the C2 sequence.

5

C1/C2 – Any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence

10

T1 sequence – Any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence. The T1 and T2 sequences must be between about 1kb and 105kb apart.

15

T2 sequence – Any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence. The T2 and T1 sequences must be between about 1kb and 105kb apart.

20

Last exon gap or Gap-Distance – The number of bases between the end of transcription and the beginning of the C1/C2 sequence. In prokaryotes and single-celled eukaryotes this gap can range from no bases to 500 bases. In multi-celled eukaryotes the gap can be as large as 10,000 bases.

Poly-adenylation signal – A number of Adenosine (A) bases are added to the mRNA at the end of the 3'UTR.

25

Possible Connectron – Any set of T1, T2 and C1/C2 sequences such that the C1 sequence is identical to the T1 sequence and the C2 sequence is identical to the T2 sequence. The promoter of some gene causes the mRNA of the gene to be expressed. The mRNA is edited to eliminate the introns. The whole mRNA including the 3'UTR can move about in the cell or the nucleus of the cell. The C1/C2 RNA that is part of the 3'UTR moves to the T1 and T2 DNA sequences. A triple-stranded complex of the DNA and the RNA forms such that the C1 sequence forms hydrogen bonds with the T1 sequence and the C2 sequence forms hydrogen bonds with the T2 sequence.

30

Because the C1 sequence is adjacent to the C2 sequence, the T1 sequence is brought physically close to the T2 sequence. This produces a loop of between about 1kb and 105kb in the DNA. Histone proteins reduce the length of the DNA by binding 200 bases. Histone/DNA complexes form six-fold symmetry chromatin assemblies. The diameter of the chromatin assemblies is approximately 30nm.

Real Connectron – Any Possible Connectron which is within the Gap-Distance of some gene

Homologous connectron – The T1 sequence and the T2 sequence are on the same chromosome as the C1/C2 sequence

Heterologous connectron – The T1 sequence and the T2 sequence are on a chromosome different from chromosome of the C1/C2 sequence

Permanent connectron – Any C1/C2 sequence, which is 3' UTR to some gene that is not surrounded by any T1 and T2 sequence pairs

Transient connectron – Any C1/C2 sequence, which is 3' UTR to some gene that is surrounded by one or more T1 and T2 sequence pairs

Self-limiting connectron – Any C1/C2 sequence which is 3'UTR to some gene that is surrounded by the T1 and T2 sequences such that C1=T1 and C2=T2

Geneless connectron - Any C1/C2 sequence which is not 3'UTR to some gene but is surrounded by some T1 and T2. A promoter may lie 5' to the C1/C2 sequence.

Bidirectionality of Connectron Excitation – A C1/C2 short loop on one strand selects a T1-T2 long loop pair on the same or the opposite strand. The C1/C2 short loop has a complementary C1'/C2' sequence on the opposite strand. Similarly the T1-T2 long loop pair has a complementary long loop pair T1'-T2'. Wherever a C1/C2, T1-T2 tetrad exists there is a complementary C1'/C2', T1'-T2' tetrad. The C1/C2 short loop

5

T1 T2 gene - C1/C2

+ strand -----

- strand -----

10

T1 T2

+ strand -----

- strand -----

C2/C1 - gene

15

+ strand -----

- strand -----

T2' T1' C2'/C1' - gene

20

gene - C1'/C2'

+ strand -----

- strand -----

T2' T1'

Of course, the short loops and the long loops do not have to be on the same chromosome.

25 Hierarchy of connectron action – When a C1/C2 is expressed it forms a T1-T2 loop by forming a connectron. The C1/C2 sequence does not have to be on the same chromosome as the T1 and T2 sequences. This provides a way of causing interaction between chromosomes. When the T1-T2 loop forms, any genes in that loop region which had been expressing C1/C2 sequences in their 3'UTRs, now cease expressing

30 the C1/C2 sequences. The connectrons formed by these C1/C2 sequences will cease to exist after some time thus opening up the genes inside the respective T1-T2 loops

to expression. The hierarchy of connectron action is alternates between repression and expression. The connectron hierarchies can be of any depth.

5 One-to-Many connectron action – One C1/C2 sequence can form connectrons in many different places on many different chromosomes. The only requirement is that C1=T1 and C2=T2. This makes it possible for one expression event to control the expression of many genes on different chromosomes.

10 Many-to-One connectron action – C1/C2s that come from many different places on many different chromosomes can form a connectron for a specific T1-T2 sequence pair. The only requirement is that C1=T1 and C2=T2. This makes it possible for many different expression events to control the expression of one set of genes on a particular chromosome.

15 Many-to-Many connectron action – The arrangement of C1/C2s and T1-T2s across chromosomes can form a complex web of gene expression control relationships.

20 Percentage of the Genome Regulated by Connectrons – Since the connectrons for a sequenced genome can be calculated, the percentage of the genome that is open to connectron regulation can be known.

25 Emergent Property – The network of connectrons in any genome emerges from a knowledge of the complete DNA sequence of the genome. Because both the C1/C2 sequences and the T1-T2 sequences can be any place in the genome, the whole genomic sequence must be known before all the connectrons can be determined.

30 Paradigm Shift – For the past fifty years since the discovery by Watson and Crick of the double-helical nature of DNA, the reigning paradigm for scientific discovery has been the study of one gene and its effects on the behavior of a cell. The advent of genomic sequencing and this invention of connectrons that emerge from the whole genome will produce a shift in the way scientists view biological systems and the way they formulate and execute experiments. The many-to-many relationships between

the connectrons means that there are many ways in which the expression of a set of genes can be modulated. The multiplicity of control pathways means produces a system stability that makes it possible for biological systems to be stable for long periods of evolutionary time. The thinking that goes into formulating scientific experiments will have to change to accommodate the changes in understanding that will be induced by the application and extension of this patent application.

Hierarchy of DNA Structuring – The DNA of a cell's genome is structured in a hierarchy of six levels. Figures 1, 2 and 3 have been adapted from The Molecular Biology of the Cell by Alberts, Bray, Lewis, Raff, Roberts and Watson [third edition pages 354, 345 and 348]. As shown in figure 1, the double stranded DNA is level 1. The double-stranded DNA is wrapped around histone proteins to form a chromatin particle that is level 2 of the hierarchy. Level 2 is described as “beads-on-a-string” in figure 1. The chromatin particles are packed in a six-fold symmetry as shown in figure 2a and figure 2b. These six-fold assemblies have a diameter of 30 nm. Each 30 nm assembly contains from 18 (i.e. $6 * 3$) to 30 (i.e. $6 * 5$) chromatin particles. The 30 nm assemblies aggregate into large loops which range in length from 5,000 bases to 100,000 bases of DNA. The size of these large loops as shown in figure 1 is approximately 300 nm. These large loops constitute level 4 of the structuring hierarchy. As shown in figure 1, level 5 of the DNA structuring hierarchy many large loops are condensed to form a structure which is approximately 700 nm in diameter. The complete chromosome that constitutes level 6 of the hierarchy is composed of two very long sections of level 5 DNA.

Model of Chromatin Structure – The level 4 structure of DNA as shown in figure 1 ranges in length from 5,000 to 105,000 bases of DNA. Figure 3 shows that proteins are thought to connect portions of the long loops formed by the 30 nm particles to form a chromosome axis. These condensed long loops are described as chromomeres in The Molecular Biology of the Cell.

Prior Art

The chromomere model of DNA structuring was presented by N. A Resnik, et al.[1] and is based on electron microscopic data. There are more recent papers studying a variety of genomes with electron microscopy but no equivalent study of chromomeres has been done on a fully sequenced genome.

A recent News Feature in Nature by T. Gura [2] described the discovery of post-transcriptional gene silencing in which viral RNA interacts with the transcribed RNA of the cell to silence the expression of genes. This article describes experiments in *C. elegans* and *D. megalomaster* in which RNA that is complementary to mRNA introduced into a cell. This “antisense” RNA has the effect of turning off the expression of one or more genes. The introduced complementary RNA produces an “RNA interference” called RNAi.

Thomas Werner and his colleagues at Genomatix in Munich, Germany have developed an approach to understanding what they call “Matrix Attachment Region” (MAR). Figure 5 shows their interpretation of the structure of DNA surrounding a gene. The following description of the MAR is copied from the Genomatix web site

“Matrix Attachment Regions (MARs) MARs are sequence regions that are responsible for the attachment of genomic DNA to the nuclear matrix or scaffold. Transcription absolutely requires anchorage of genomic DNA to the nuclear matrix.

Functional features of MARs:

Anchoring of regulatory elements like promoters and enhancers to the nuclear matrix.

Ensuring long term activity of promoters and enhancers in chromatin.

Insulation, rendering a functional domain insensitive to position effects.

Genomatix is conducting a research project to define and detect MARs by computer-analysis.”

0986925 053004
T00250 52599850

Brief Description of the Objects of the Invention

5 An object of the invention is to provide a method of identifying DNA sequences that control the expression of different collections of genes in a genome comprising, detecting selected DNA sequences adjacent to some genes excluding exons and introns.

10 An object of the invention is to provide a method of identifying DNA sequences that control the expression of different collections of genes comprising, detecting, by computer, one or more pairs of non-adjacent DNA sequences to which are bound to two RNA sequences.

15 An object of the invention is to provide a method of identifying DNA sequences that control the expression of different collections of genes in a genome comprising detecting changes in connectron behavior in the genome.

20 An object of the invention is to provide a method of modifying the expression of different gene collections in a genome, comprising detecting changes in connectron behavior as a result of an exogenous stimulus.

25 An object of the invention is to provide a method of detecting where and when new genes are being integrated into a host genome comprising detecting the connectrons in said host genome.

 An object of the invention is to provide a method of detecting the expression effect of different gene collections in a given body comprising detecting the back and forth flow of connectrons between the chromosomes thereof.

An object of the invention is to provide a method of modifying a given body comprising modifying the connectron organization therein.

An object of the invention is to provide a method of detecting connectron control and target sequences in a given genome comprising:

determining the base composition of said genome,
determining one or more sites of control sequence organization, and/or
determining one or more sites of target application.

An object of the invention is to provide a method of determining the response of a cell in any tissue to changes in the cell's environment and/or genetic composition comprising providing a complete genomic DNA sequence for the organism and determining the effect of changes in connectrons due to application of a given exogenous stimulus to the genome.

An object of the invention is to provide a method of determining in prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes, the tetradic relationship $T1=C1$ and $T2=C2$ where T1 and T2 are DNA sequences 20 or more bases in length, where the C1 sequence is adjacent to the C2 sequence, where the T1 and T2 sequences are on the same chromosome, and where the C1/C2 sequences are on the same chromosome as T1 and T2 or where the C1/C2 sequences are on a chromosome different from T1 and T2, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

An object of the invention is to provide a method of determining in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits many different C1/C2 short loops to control the existence of a T1-T2 long loop and wherein said C1/C2 short loops can be on the same chromosome or on different chromosomes from the T1-T2 long loop, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

5 An object of the invention is to provide a method of determining in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of many T1-T2 long loops, the C1/C2 short loop can be on the same chromosome or on different chromosomes from the T1-T2 long loops, wherein:

10

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

15

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

20

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

25

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

30

An object of the invention is to provide a method of determining in the connectron relationships between prokaryotes and their plasmids wherein said connectrons implement a control mechanism between the two genomes that makes it possible from

them to form a symbiotic relationship, and in the case of *D. radiodurans* the relationship is not symmetric, and the *D. radiodurans* genome sends C1/C2 short loops to the MP1 plasmid, wherein:

5 C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

10 C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

15 C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

20 T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

25 T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

30 An object of the invention is to provide a method of determining that connectron relationships that exist in plant and higher animals.

An object of the invention is to provide a method of determining in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of one or more T1-T2 long loops without being subject to any expression controls other than those of the gene to which the C1/C2 is 3'UTR, wherein:

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

5 C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

10 C1/C2 - any positive or negative strand DNA sequence of 540 or more bases such that the C1 sequence is adjacent to the C2 sequence,

15 T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart,

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart, and

20 3'UTR - untranslated 3' end of an mRNA is beyond the end of the last exon, a stop codon in the mRNA causes the ribosome to stop the translation of mRNA into protein.

25 An object of the invention is to provide a method of determining in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of one or more T1-T2 long loops such that this C1/C2 short loop is itself subject to expression control by another T1-T2 long loop which surrounds it, wherein:

30 C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

5

C1/C2 - any positive or negative strand DNA sequence of 540 or more bases such that the C1 sequence is adjacent to the C2 sequence,

10

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

15

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

An object of the invention is to provide a method of determining in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes, the connectron relationship that permits one C1/C2 short loop to control the existence of the T1-T2 long loop that surrounds it, wherein:

20

C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

25

C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

30

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

5 T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

10 An object of the invention is to provide a method of determining the connectron relationships that do not have any genes within the T1-T2 long loop, wherein:

T1 sequence is any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, and

15 T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, and the T2 or T1 sequences must be between about 1kb and 105kb apart.

20 An object of the invention is to provide a method of determining the geneless connectron relationship where one C1/C2 short loop controls the existence of many geneless T1-T2 long loops, wherein:

25 C1 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C2 sequence must occur in the same chromosome as the C1 sequence,

30 C2 sequence - any positive or negative strand DNA sequence of 20 bases or more, the C1 sequence must occur in the same chromosome as the C2 sequence,

C1/C2 - any positive or negative strand DNA sequence of 40 or more bases such that the C1 sequence is adjacent to the C2 sequence,

5

T1 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T2 sequence, the T1 and T2 sequences must be between about 1kb and 105kb apart, and

T2 sequence - any positive or negative strand DNA sequence of 20 bases or more that is on the same chromosome as the T1 sequence, the T2 or T1 sequences must be between about 1kb and 105kb apart.

09866925:053001
T00E50:52699860

Description of the Drawings and Tables

The above and other objects, advantages and features of the invention will become more apparent when considered with the following specification and accompanying drawings and tables wherein:

Figure 1 DNA is structured in six levels of increasing condensation. Double stranded DNA is level 1. Two turns of DNA are wrapped about each chromatin particle at level 2. The chromatin particles which each containing 200 base pairs form into 30 nm particles at level 3. The 30 nm particles form into large loops with an approximate dimension of 300 nm at level 4. Metaphase chromosomes form a condensed structure with an approximate dimension of 700 nm at level 5. An entire metaphase chromosome has a width of approximately 1400 nm at level 6. The large loops at level 4 of the DNA structuring are thought to have between 20,000 (20 kb) and 100,000 (100 kb) base pairs.

The Molecular Biology of the Cell by Alberts, Bray, Lewis, Raff, Roberts and Watson, 3rd. ed. , Garland Publishing, Inc., New York, 1994, p. 354

Figure 2 (a) Chromatin DNA forms into a six-fold symmetry 30nm particles.
(b) The six-fold symmetry 30nm particles form a linear chain with a varying number of repeat units.

The Molecular Biology of the Cell by Alberts, Bray, Lewis, Raff, Roberts and Watson , 3rd. ed. , Garland Publishing, Inc., New York, 1994, p. 345

Figure 3 Long loops of 30nm particles are thought to be closed at the bottom of the loop by proteins.

5 The Molecular Biology of the Cell by Alberts, Bray, Lewis, Raff, Roberts and Watson, 3rd. ed. , Garland Publishing, Inc., New York, 1994, p. 348

Figure 4 (a) Transcription and Editing. (b) Movement of the RNA through the Nucleus. (c) Connectron Formation

10 Figure 5 Overview of schematic organization of a typical transcriptionally active chromosomal loop.
From http://genomatix.gsf.de/func_genomics/functional_genomics.html

15 Table 1 Connectron Properties for Prokaryotic, Archea and Eukaryotic Genomes

Table 2 Yeast Inter-Chromosomal Connectron Distribution

100250 5269960

	Figure 6	Genome size plotted as a log-log function of the Number of Connectrons
	Figure 7	Number of Sequence Instances plotted as a function of the Number of Fragments
5	Figure 8	Level 0 – The overall view of the algorithm
	Figure 9	Level 1 – Process Flow of the Algorithm
	Figure 10	Level 2a – two pages – Process Genome into Blocking Fragment File
	Figure 11	Level 2b – two pages – Compute the Connectrons for a Genome
	Figure 12	Level 2c – two pages – Analyze Possible Connectrons
10	Figure 13	Level 3a – Setup Genome Usage Memory
	Figure 14	Level 3b – Find DBP-Size Blocking File for T1-Window
	Figure 15	Level 1 – Find DBP-Size Blocking File for T2-Window
	Figure 16	Level 2a – two pages – Find C1/C2 Entries
15	Figure 17	Level 2b – two pages – Scan Genome Usage Memory for Potential Connectrons

Description of the Invention

A connectron is a relationship among four DNA sequences. Each sequence must be at least 20 bases long. There is a report by Sharp and Zamore [3] that RNA sequences of “about length 25” are important as sources of RNAi. 27 bases were actually used as the minimum length of each of the sequences. The T1 sequence is on one strand of some chromosome in a genome. The T2 sequence is on the same strand of the same chromosome as the T1 sequence. The T1 and T2 sequences (which are each at least 20 bases in length) must be at least 5,000 bases distant from each other but they can not be more than 105,000 bases distant from each other. The C1 sequence and the C2 sequence (which are each at least 20 bases in length) are adjacent to each other on some strand of some chromosome in the genome. The C1/C2 sequences – called the “short loop” – can be on the same strand as the T1 and T2 sequences or they can be on the opposite strand. The C1/C2 sequences of the short loop can be on the same chromosome as the T1 and T2 sequences but they can also be on a different chromosome in the genome. When a genome has only one chromosome, then the point is moot. Many genomes, of course, have several chromosomes. The C1 sequence is identical to the T1 sequence and the C2 sequence is identical to the T2 sequence.

The C1/C2 sequence must be on the same strand as a gene, either be directly adjacent to the gene (i.e. a gap of 0 bases) for prokaryotic genomes or at this time be within 10,000 bases for eukaryotic genomes. The size of the gap between the end of the gene and the beginning of the C1/C2 sequence is a variable. The C1/C2 short loop is expressed as the 3'UTR (Un-Translated Region) of the gene. In the case of prokaryotic genes that do not normally have introns, the whole mRNA becomes the active species for connectron formation. In the case of eukaryotic genes, the whole transcript is the active species for connectron formation upon editing of the transcript to eliminate the introns. The whole transcript then can move about in the cytoplasm of prokaryotic cells or the nucleus of eukaryotic cells. Since the C1 sequence is equivalent to the T1 sequence and the C2 sequence is equivalent to the T2 sequence,

the C1 RNA can form a Hoogsteen triple-stranded RNA/DNA/DNA helix with the double-stranded T1 sequence. Similarly the C2 RNA can form a Hoogsteen triple-stranded RNA/DNA/DNA helix with the double-stranded T2 sequence. Because the C1 sequence and the C2 sequence are adjacent to each other, the C1/T2 RNA/DNA/DNA Hoogsteen triple helix is brought into physical adjacency to the C2/T2 RNA/DNA/DNA Hoogsteen triple helix. RNA/DNA/DNA hybrid helices are the most stable form of triple helix. RNA double helices, DNA double helices, RNA triple helices and DNA triple helices are all significantly less stable than a RNA/double-stranded DNA triple helix. The stable physical adjacency of the two triple-stranded Hoogsteen helices ensures that the long loop of double-stranded DNA between the T1 sequence and the T2 sequence can then be structured into 30 nm chromatin particles as shown in level 4 of figure 1. The genes on either strand of the DNA between the T1 sequence and the T2 sequence when they are structured into the 30 nm chromatin particles are not open to promotion and expression.

The tetradic relationship between the T1 and T2 sequences that form the long loop and the C1/C2 sequences that form the short loop are called a connectron. The name “connectron” was suggested by J. David Rawn Ph.D. of Towson University. A connectron is possible if the T1, T2, C1 and C2 sequences exist. A connectron is real if the C1/C2 short loop sequence is adjacent to an expressible gene. If the expression of the adjacent gene is inside one or more T1 – T2 long loops then this connectron is said to be transient. If the adjacent gene is not inside any possible T1-T2 long loop then the connectron is said to be permanent. If a connectron is inside of a T1-T2 long loop that has the same sequences (i.e. T1 is really equal to C1 and T2 is really equal to C2) then the connectron is said to be self-limiting. This is true because once the C1/C2 sequence is expressed it forms the T1-T2 long loop that then shuts off the expression of the gene adjacent to the C1/C2 sequence. Self-limiting connectrons can also be called “spike” connectrons since they generate a short-duration spike of the C1/C2 short loop sequence. If a T1-T2 long loop does not contain any genes but it contains C1/C2 short loop sequences then this type of connectrons is said to be geneless. The C1/C2 short loops within a geneless T1-T2 long loop can, of course, control the expression of genes.

5 The physical existence and lifetimes of the connectrons must be proved by molecular biological experimentation. This physical experimental process, however, is logically quite separate from the computational experimentation that have been conducted from June of 1999 to May of 2001. The computational search for the existence of connectrons has been extremely positive. These computations have shown that connectrons exist in prokaryotes, in archea, between prokaryotes and their plasmids, in single-celled eukaryotes, in multi-celled eukaryotes, in plants, in higher animals and in humans. All of these features and properties are described in the claims section that follows.

10 The connectron invention is very powerful. It depends only on sequence equivalency. The minimum length of the four sequences seems to be about 20 bases. In the calculations shown in this patent application, 27 bases have been used as a minimum. The Nature News Feature [1] says that other scientists have found RNA sequences of length about 25 that have interesting gene silencing properties. The Nature article does not give any mechanism. Because of my algorithm and its use on a variety of genomes, this patent application provides the computational proof that a particular mechanism is highly probable. The connectron invention provides an explanation for how communication occurs with a chromosome as well as between chromosomes in genomes that have more than one chromosome. Since each T1-T2 long loop can contain one or more genes, the connectron invention provides a mechanism for turning on and turning off sets of genes simultaneously. In time, the connectron invention will provide an explanation for how differentiation of how one cell's behavior differs from the behavior of another adjacent cell. It is already clear from the computational experiments that have been made on *S. cervesiae*, *C. elegans* and *D. megalomaster* that the number of geneless connectrons increases dramatically as evolution proceeds from single-celled eukaryotes (i.e. *S. cervesiae*) to 1,000 cell eukaryotes (i.e. *C. elegans*) to visible creatures (i.e. *D. megalomaster*). The extension of this evolutionary progress to plants (i.e. *A. thaliana*) for which only three chromosomes are sequenced and humans (i.e. *H. sapiens*) for which only one chromosome is completely sequenced. Although the complete human genome was

published in Nature and Science in February of 2001, the NIH-sponsored genomic sequencing results are available for about 1/3 of the bases in the whole genome. The human genomic sequence determined by Celera Genomics, Inc. is available only by subscription. Table 1 shows how the genome size, the number of genes, the number of gene-containing and geneless connectrons and the percentage of genes controlled are related in many different genomes.

The C1/C2 short loops originate on one chromosome. The T1-T2 long loops can be on the same or different chromosomes. Table 2 which is for yeast (*S. cerevisiae*) is a square matrix of how many C1/C2 short loops on a given chromosome are sent to form T1-T2 long loops on other chromosomes. The diagonal of this matrix shows that many chromosomes send connectrons to themselves. The striking feature of this particular table is that chromosome 6 only sends connectrons to chromosome 12 but that it receives connectrons from chromosomes 4,5,7,10,12,13,15 and 16.

Any tetrad of connectron sequences (i.e. the T1, T2, C1 and C2 sequences) as well as the fact of the adjacency of the C1/C2 short loop sequence to the transcribing gene can be patented because the content of matter and the utility can be exactly described. The utility of a connectron is that the T1-T2 long loop shuts off the expression of the genes that lie between the T1 sequence and the T2 sequence. In the case of geneless connectrons, the utility is of a higher level in that the C1/C2 short loops contained in the higher-level geneless T1-T2 long loop, eventually form other lower-level T1-T2 long loops around a set of genes.

The invention of connectrons comes at a particularly important time in biological discovery. The geneless connectrons make a many-to-many hierarchical control mechanism possible. It is already clear from the determination of the connectrons for *C. elegans* and *D. megalomaster* that there are as many or more geneless connectrons than there are genes. It has been clear for some time that the number of genes in a genome is not particularly correlated with the size of the genome. Figure 6 shows that the size of a genome is roughly linearly correlated with the number of connectrons.

The connectron invention can be used to generate a model of behavior in any cell. The simulation of connectron behavior in different genomes will be the subject of another patent application.

5

The connectron invention provides for a rational exploitation of the information contained in the raw genomic DNA sequence by forming a hierarchy of relationships between geneless connectrons, transient connectrons, permanent connectrons, self-limiting connectrons and the expression of genes.

10

0966923 "053001
"0050" 5269990

Detailed Description of the Invention

The algorithm for the determination of connectrons in any genome or any genome fragment is represented in the following flow diagrams. The Level 0 diagram in figure 8 shows the general relationships in a digital computer. The central processor of the digital computer uses the computer program to take genome descriptors, the genomic DNA sequences and the tables of gene features to produce a file of blocking fragments and a file of the optimal connectrons for the genome. The printer serves to make hard copies of the files and this patent application. The level 1 diagram in figure 9 shows the three essential steps in the determination of connectrons. The genome is first processed into a blocking fragment file. Then the blocking fragments are used to compute the connectrons for the genome. Finally the potential connectrons are analyzed to determine if the C1/C2 sequences are in the 3'UTR of a gene. The level 2a diagram in figure 10 shows the steps required for the processing of the genome into a file of blocking fragments. The genomic DNA sequence is decomposed into 27-base frames for both the positive and negative strands. These fragments are written to the unsorted fragment file. The fragment file is then sorted is then read and formed into groups of equivalent sequences. The (.blk) file contains the sequence and a pointer to the (.gptr) file which contains the pointers to the position of the fragments in the genomes. The position in the genome includes the chromosome number, the position in the chromosome and the strand (i.e. positive and negative). A sample of these files follows

Sample of the (.blk) file for *S. cerevisiae*

27-base fragment Number Pointer
 of instances to (.gptr) file

11111111111111111111111111111111	0	1
111111123244233313332443414	1	2
111111141113443133314333341	2	4
111111232442333133324434141	1	5
111111323311133323144423444	2	7
111111332213331341414443413	2	9
111111333444112343412323243	1	10

5	111111333444113343412323243	9	19
	111111411134431333143333414	2	21
	111111443223134142124434124	2	23
	111112223234344444443144442	2	25
	111112244123441122214421213	8	33
10	111112311241114344334134431	2	35
	111112324423331333244341414	1	36
	111112344232231344242234342	1	37
	111112433444244421144134211	1	38
	111112444311313442332142224	1	39
15	111113131241131114424413231	1	40
	111113143332344311113133411	1	41
	111113233111333231444234441	2	43

In fragments above 1=G, 2=C, 3=A, 4=T

Sample of the (.gptr) file for *S. cerevisiae*

There are 16 chromosomes in *S. cerevisiae*

20	Item Chromosome Position Direction in Chromosome			
	1	0	0	0
25	2	4	11137	1
	3	12	467619	1
	4	12	458482	1
	5	4	11138	1
	6	12	465759	2
30	7	12	456622	1
	8	1	219366	1
	9	8	539978	1
	10	14	522451	1
35	11	4	1099073	1
	12	4	1210003	1
	13	7	539068	1
	14	12	654136	1
	15	12	596455	1
40	16	15	121016	1
	17	15	598127	2
	18	16	847724	1
	19	16	59765	1
45	20	12	467620	1
	21	12	458483	1
	22	12	461657	1
	23	12	452520	1
	24	13	838006	1

	25	15	288270	1
	26	4	83593	1
	27	4	992867	1
5	28	6	162265	1
	29	7	845687	1
	30	10	531560	2
	31	15	282208	1
	32	16	860418	1
10	33	16	572308	1
	34	12	465992	1
	35	12	456855	1
	36	4	11139	1
	37	8	89343	1
	38	4	10302	1
15	39	1	19894	2
	40	16	9311	1
	41	10	735203	1
	42	12	465760	1
20	43	12	456623	1

In direction column above 1=positive strand, 2=negative strand

The level 2b diagram in figure 11 shows the computation of the connectrons. The genome descriptors consist of the number and length of the chromosomes. The algorithm uses an array that represents several facts about each base position in the genome. The level 3a diagram in figure 13 shows the setup of the Genome-Usage memory. The gene features are used to prevent the region of the genome that codes for proteins from being used for the connectron sequences (i.e. the T1s, the T2s, the C1s and the C2s). In the level 2a diagram of figure 10, the algorithm steps through each chromosome and within each chromosome through each base position looking for acceptable T1-windows of 27 bases. A T1-window can be used to form a connectron relationship if there are two or more instances of this fragment in the blocking fragment file. The computation in the level 3b diagram of figure 14 determines if the T1-window is acceptable or not. Once an acceptable T1-window is found, the algorithm (in the level 2a diagram of figure 10) looks for acceptable T2-window positions that lie between 5,000 and 105,000 bases from the T1-window. The computation for determining acceptable T2-window positions is done in the level 3c diagram of figure 15. Once a pair of T1 and T2 window positions are found, the

algorithm looks among the instances of these T1 and T2 sequences for a pair of sequences C1 and C2 that lie within 200 bases of each other on the same chromosome. The computation for determining acceptable C1/C2 windows is shown in the level 3d diagram in figure 16. In the level 3e diagram of figure 17 the Genome-Usage memory is scanned for the Possible-Connectrons. In the level 2c diagram of figure 12 the Possible-Connectrons are scanned to determine if the C1/C2 sequences are within the Gap-Distance of a gene on either the positive or the negative strands. The Real-Connectrons are then written out in several different files including the descriptions in the claims section.

5

10

T00E50" 22699860

Examples

The algorithm for the determination of optimal connectrons has been applied to a number of different publicly available genomes. The connectron is a tetradic relationship between four sequence elements – T1, T2, C1 and C2. The claims presented in this section are written by the program NearGene that implements the flow diagram Level 2c of figure 12. The examples are written a uniform type of English. Each example contains some or all of the following elements

10 Name of genome
 Description of T1
 Length of T1-T2 loop
 The chromosome on which the T1-T2 loop exists
 The identifier number within the genome of the T1 sequence
15 The T1 sequence
 Description of T2
 The identifier number within the genome of the T2 sequence
 The T2 sequence
 A list of genes whose expression is controlled by the T1-T2 loop
20 The common names of the genes as obtained from the NCBI gene feature file
 (.ptt)
 A list of C1/C2 short loops whose expression if controlled by the T1-T2 loop
 The chromosome on which the C1/C2 short loop exists
 The common name of the gene which expresses the C1/C2 short loop as an
25 RNA
 The sequence of the C1/C2 short loop
 A list of C1/C2 short loops that control the formation of the T1-T2 loop
 The chromosome on which the C1/C2 short loop exists
 The common name of the gene which expresses the C1/C2 short loop as an
30 RNA
 The sequence of the C1/C2 short loop

The match between the C1/C2 sequence and the T1 sequence
The match between the C1/C2 sequence and the T2 sequence

5 The uniform descriptions make it possible to rapidly comprehend the specifics in each example.

When a sequence element is very long a series of four dots has been inserted between the beginning and ending sequence groups. A variable number of bases have been deleted.

10

0986925.053001
T00E50.32699860

Index of Pages for Connectron Samples

Page 39

5

1. Connectrons occur in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Page 57

10

2. Many Connectrons control the expression of one set of genes in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Page 83

15

3. One connectron controls the expression of many sets of genes in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Page 107

4. Connectrons occur between prokaryotes and their plasmids.

Page 117

20

5. Connectrons occur in plants and higher animals

Page 126

6. Permanent connectrons exist in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

25

Page 135

7. Transient connectrons exist in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Page 152

30

8. Self-limiting connectrons occur in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes

Page 164

9. Geneless connectrons exist in single-celled and multi-celled eukaryotes

5

Page 174

10. One connectron controls many geneless connectrons in single-celled and multi-celled eukaryotes

Connectron control elements for chromosome 1 of the E. coli genome

A double stranded DNA loop of length 93.542 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3197. This T1 control element has the DNA sequence

AAAAAATGCGCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3308. This T2 control element has the DNA sequence

TAAATTCCTCTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
ACACGGAACAACGGCAAACACGCCCGGGTCAGCGGGGTTCTCCTGAG
AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
GCGTATTATGCACACCCCGCGCCGCT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

rrsC	gltU	rrlC	rrfC	aspT	trpT	yifA	yifE	yifB
ilvL	ilvG_1	ilvM	ilvE	ilvD	ilvA	ilvY	ilvC	ppiC
b3776	rep	gppA	rhlB	trxA	rhoL	rho	rfe	wzzE
wecB	rffH	wecD	wecE	wzxE	yifM_2	wecG	yifK	
argX	hisR	leuT	proM	aslB	aslA	hemY	hemX	
hemD	cyaA	cyaY	b3808	dapF	uvrD	b3814	corA	
yigF	yigG	rarD	yigI	pldA	recQ	yigJ	yigK	pldB
yigL	yigM	metR	metE	ysgA	udp	yigN	ubiE	yigP

b3836 yigU yigW_1 rfaH yigC ubiB fadA fadB
 pepQ trkH hemG

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 1 whose identifier is 3204 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrsC and has the DNA sequence

GATGTGCCCAGATGGGATTAGCTAGTAGGTGGGGTAACGGCTCACCTAGG
 CGACGATCCCTAGCTGGTCTGAGAGGATGACCAGCCACACTGGAAGT
 ACACGGTCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATG
 GGCGCAAGCCTGATGCAGCCATGCCGCGTGTATGAA

A C1/C2 short loop on chromosome 1 whose identifier is 3206 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrsC and has the DNA sequence

GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
 GGGTTCGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC
 ACCTGCCTTAATATCTCAAACTCATCTTCGGGTGATGTTTGAGATATTG
 CTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAA

A C1/C2 short loop on chromosome 1 whose identifier is 3223 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrlC and has the DNA sequence

GCTGAAGTAGGTCCCAAGGGTATGGCTGTTGCGCCATTTAAAGTGGTACGC
GAGCTGGGTTTAGAACGTCGTGAGACAGTTCGGTCCCTATCTGCCGTGGG
CGCTGGAGAACTGAGGGGGGCTGCTCCTAGTACGAGAGGACCGGAGTGG
ACGCATCACTGGTGTTCGGGTTGTCATGCCAATGGCA

5

A C1/C2 short loop on chromosome 1 whose identifier is 3225 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrlC and has the DNA sequence

10

AAACAGAATTTGCCTGGCGGCCGTAGCGCGGTGGTCCCACCTGACCCCAT
GCCGAATCAGAAGTGAAACGCCGTAGCGCCGATGGTAGTGTGGGGTCTC
CCCATGCGAGAGTAGGGAAGTGCCAGGCATCAAATTAAGCAGTA

15

A C1/C2 short loop on chromosome 1 whose identifier is 3228 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrfC and has the DNA sequence

20

GGTCATAAAACCGGTGGTTGTAAAAGAATTCGGTGGAGCGGTAGTTCAGT
CGGTTAGAATACCTGCCTGTCACGCAGGGGGTCGCGGGTTCGAGTCCCGT
CCGTTCCGCCAC

25

A C1/C2 short loop on chromosome 1 whose identifier is 3301 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ubiB and has the DNA sequence

30

TTATCGTGCCTACAAATAGTCCGAACCGTAGGCCGGATAAGGCGTTTACG
CCGCATC

A C1/C2 short loop on chromosome 1 whose identifier is 3307 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene fadA and has the DNA sequence

5

TGCCGGATGCGGCGTAAACGCCTTATCCGGCCTACGGTTCGGACTATTTGT
AGGCA

10 A C1/C2 short loop on chromosome 1 whose identifier is 3327 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene hemG and has the DNA sequence

15 AAAAAATGCGCGGTCAGAAAATTATTTTAAATTTCTCTTGTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAAGGCGTATTATG...CCCGTCACACCA
TGGGAGTGGGTTGCAAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCT
20 TACCACTTTGTGATTCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTA
GGGGAACCTGCGGTTGGATCACCTCCTTACCTTAAAGAAGCGTTCTTTG

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 1 whose identifier is 3307 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene hemG and has the DNA sequence

30 AAAAAATGCGCGGTCAGAAAATTATTTTAAATTTCTCTTGTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAAGGCGTATTATG...CCCGTCACACCA

TGGGAGTGGGTTGCAAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCT
TACCACTTTGTGATTCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTA
GGGGAACCTGCGGTTGGATCACCTCCTTACCTTAAAGAAGCGTTCTTTG

5 The match between the T1 sequence and the C1/C2 sequence is

AAAAAATGCGCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAA

The match between the T2 sequence and the C1/C2 sequence is

TAAATTCCTCTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
ACACGGAACAACGGCAAACACGCCCGGGTCAGCGGGGTTCTCCTGAG
AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
GCGTATTATGCACACCCCGCGCCGCT

A C1/C2 short loop on chromosome 1 whose identifier is 3432 controls the
expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
as a RNA single strand that is 3'UTR to the gene btuB and has the DNA sequence

TGCGCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGGAATAACT
CCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACGCCGCCGG
GTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAAATAAATG
CTTGACTCTGTAGCGGGAAGGCGTATTATGCACACC...ACACCATGGGAGT
GGGTTGCAAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCTTACCACT
TTGTGATTCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTAGGGGAAC
CTGCGGTTGGATCACCTCCTTACCTTAAAGAAGCGT

The match between the T1 sequence and the C1/C2 sequence is

TGCGCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGGAATAACT
CCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACGCCGCCGG
GTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAAATAAATG
CTTGACTCTGTAGCGGGAA

5

The match between the T2 sequence and the C1/C2 sequence is

TAAATTCCTCTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
ACACGGAACAACGGCAAACACGCCGCCGGGTCAGCGGGGTTCTCCTGAG
AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
GCGTATTATGCACACCCCGCGCCGCT

10

A C1/C2 short loop on chromosome 1 whose identifier is 2218 controls the
expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
as a RNA single strand that is 3'UTR to the gene clpB and has the DNA sequence

15

CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
AACGGCAAACACGCCGCCGGGC

20

The match between the T1 sequence and the C1/C2 sequence is

CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
AACGGCAAACACGCCGCCGGGC

25


The match between the T2 sequence and the C1/C2 sequence is

CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
AACGGCAAACACGCCGCCGGGTC

30

Example of an archea connectron – *H. pylori*

In this example the existence of the T1-T2 (812-882) long loop is controlled by three C1/C2 short loops (881, 813 and 1241). The T1-T2 long loop controls the expression of 54 genes on chromosome 1 in addition to one C1/C2 (843) short loop.



Connectron control elements for chromosome 1 of *H. pylori* genome

A double stranded DNA loop of length 96.385 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 812. This T1 control element has the DNA sequence

TTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 882. This T2 control element has the DNA sequence

TAGCGGAACTAAAGCATTCATCCCAAACACTAAAGATATTTGG

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

HP0999	HP1000	HP1001	HP1002	HP1003	HP1005	HP1006
HP1008	HP1009	HPtRNA-Pro	HP1010	HP1011	HP1013	HP1015
HP1017	HP1018	HP1020	HP1021	HP1022	HP1023	HP1024

	HP1025	HP1027	HP1028	HP1030	HP1031	HP1033	HP1034
	HP1038	HP1039	HP1040	HP1041	HP1042	HP1043	HP1044
	HP1045	HP1046	HP1051	HP1052	HP1055	HP1056	HP1058
	HP1060	HP1065	HPtRNA-Ser	HP1066	HP1067	HP1069	HP1070
5	HP1074	HP1075	HP1076	HP1077	HP1078	HP1079	HP1080
	HP1081	HP1083	HP1084	HP1085	HP1088	HP1091	HP1092
	HP1093	HP1094	HP1095	HP1096			

10 This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15 A C1/C2 short loop on chromosome 1 whose identifier is 813 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP0998 and has the DNA sequence

TTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
AAACACTAAAGATATTTGG

20 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 1 whose identifier is 881 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP1096 and has the DNA sequence

TTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
AAACACTAAAGATATTTGG

30 The match between the T1 sequence and the C1/C2 sequence is

TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCA

The match between the T2 sequence and the C1/C2 sequence is

5 TAGCGGAACTAAAGCATTTCATCCCAAACACTAAAGATATTTGG

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

10 A C1/C2 short loop on chromosome 1 whose identifier is 813 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP0998 and has the DNA sequence

15 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
AAACACTAAAGATATTTGG

A C1/C2 short loop on chromosome 1 whose identifier is 881 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP1096 and has the DNA sequence

20 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
AAACACTAAAGATATTTGG

The match between the T1 sequence and the C1/C2 sequence is

25 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCA

The match between the T2 sequence and the C1/C2 sequence is

30 TAGCGGAACTAAAGCATTTCATCCCAAACACTAAAGATATTTGG

A C1/C2 short loop on chromosome 1 whose identifier is 1241 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP1535 and has the DNA sequence

5 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
AAACA

The match between the T1 sequence and the C1/C2 sequence is


10 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCA

The match between the T2 sequence and the C1/C2 sequence is

15 TAGCGGAACTAAAGCATTTCATCCCAAACA

Example of single-celled connectron – *S. cerevisiae*

20 In this example the existence of the T1-T2 (1352-1416) long loop on chromosome 4 is controlled by one C1/C2 short loop (4213) on chromosome 10. The T1-T2 long loop controls the expression of 34 genes on chromosome 4 in addition to one C1/C2 (1356) short loop.



Connectron control elements for chromosome 1 of *S. cerevisiae* genome

A double stranded DNA loop of length 68.908 kilo-bases on chromosome 4 is bounded on the left by a T1 sequence whose identifier is 1352. This T1 control element has the DNA sequence

TTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1416. This T2 control element has the DNA sequence

ATTAGATCTATTACATTATGGGTGGTATGTTGGAATAAAAATCAACTATCA
TCTACTAACTAGTATTTACGTTACTAGTATATTATCATATACGGTGTTAGA
AGATGACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAA
ACGCAAGGATTGATAATGTAATAGGATCAATGAATATTAACATATAAAAC
GATGATAATAATATTTATAGAATTGTGTAGAATTGCAGATTCCCTTTTATG
GATTCCTAAATCCTTGAGGAGAACTTCTAGTATATCTACATACCTAATATT
ATAGCCTTAATCACAATGGAATCCCAACAATTACATCAAAATCCACATTC
TCTACAGTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

YDR170W-A	YDR171W	YDR172W	YDR173C	YDR174W	YDR175C
YDR176W	YDR177W	YDR178W	YDR179C	YDR179W-A	YDR180W
YDR181C	YDR182W	YDR183W	YDR184C	YDR185C	YDR186C
YDR187C	YDR188W	YDR189W	YDR190C	YDR191W	YDR192C
YDR193W	YDR194C	YDR195W	YDR196C	YDR197W	YDR198C
YDR199W	YDR200C	YDR201W	YDR202C	YDR203W	YDR204W
YDR205W	YDR206W	YDR207C	YDR208W	YDR209C	YDR210W

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 4 whose identifier is 1356 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YDR170W-A and has the DNA sequence

AATCACACTAATCATTCTGATGATGAACTCCCTGGACACCTCCTTCTCGAT
TCAGGAGCATCACGAACCCTTATAAGATCTGCTCATCACATACTCAGC
ATCATCTAATCCTGACATAAACGTAAGTTGATGCTCAAAAAAGAAATATAC
CAATTAACGCTATTGGTGACCTACAATTTCACTTCCAGGACAACACCAAA
ACATCAATAAAGGTATTGCACACTCCTAACATAGCCTATGACTTACTCAGT
TTGAATGAATTGGCTGCAGTAGATATCACAGCATGCTTTACCAAAAACGT
CTTAGAACG

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 10 whose identifier is 4213 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YJR029W and has the DNA sequence

ATCTATTACATTATGGGTGGTATGTTGGAATAAAAAATCCACTATCGTCTAT
CAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGATGAT
GACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAACGC
AAGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGA
ATGAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTGA
GGATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATA
TTATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACAT

The match between the T1 sequence and the C1/C2 sequence is

TTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAA

The match between the T2 sequence and the C1/C2 sequence is


5 ATCTATTACATTATGGGTGGTATGTTGGAATAAAAATC

10

Example of a multi-celled connectron – *C. elegans*

In this example the existence of the T1-T2 (9-138) long loop on chromosome 1 is controlled by three C1/C2 short loops on chromosome 5 (21719, 21949 and 21655).

15 The T1-T2 long loop controls the expression of four genes on chromosome 1 in addition to seven C1/C2 (119, 122, 125, 130, 132, 134 and 136) short loops.



30

A double stranded DNA loop of length 41.978 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 95. This T1 control element has the DNA sequence

35

0966925-053001

CAGCACGTTCTTAACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTC
CCGC

5 This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 138. This T2 control element has the DNA sequence

ACTCTGCGTCTCTTCTCCCGCATTTTTTGTAGATCA

10 This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

Y73A3A.1 Y73A3A.1 ZC123.3 ZC123.2

15 This long T1/T2 double stranded DNA loop modulates the expression of the
following C1/C2 short loops

20 A C1/C2 short loop on chromosome 1 whose identifier is 119 controls the expression
of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene ZC123.3 and has the
DNA sequence

TTGAGAACTCTGCGTCTCAACTCCCGCATTTTTTGTAGATCTACGTAGATC
AAACCGAAATGGGACACT

25 A C1/C2 short loop on chromosome 1 whose identifier is 122 controls the expression
of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene ZC123.3 and has the
DNA sequence

30 GCACGGGGTTCTGGCCTTCCTCATTGAATTTTTTCGCGCTCCATTGACAATC
GCCTGCCGGACAACGCGTGGGAAAGTCGTGTACTCCAC

A C1/C2 short loop on chromosome 1 whose identifier is 125 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.3 and has the DNA sequence

5

ACGCGCCGTAAATCTACCCCAGATATGGCCGAGCCAAAATGGCCTAGTTC
GGCAAACCTCTTTCATTTCAATTTATGAGGGAAGCCAGAA

10 A C1/C2 short loop on chromosome 1 whose identifier is 130 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.2 and has the DNA sequence

15

CTCCCGCATTTTTTGTAGATCTACGTAGATCAAACCGAAATGAGGCACTTT
CTGAATCCACGAGCTAGGCTTAAGCTTAGGCTTAAGCTTAGGCCTTTTCTC
AGGCTTAGGCTTAGGCTTA

20

A C1/C2 short loop on chromosome 1 whose identifier is 132 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.2 and has the DNA sequence

25

GCTTATGCTTGGGCTTAGGCTTAGGCGTAGGCTTAGGCTTAGGCTTAGGCT
TATGCTTAGACTTAGTCTCACTATCAGTCTTAGGCTTAGGCTTAGACTTAG
GCTTAAGCTTAGGCTTAAGCTTAGACTTAGGCTTAGGCTTAGGCTTAGGCT
TAGGCTTAGGTTTGGGCTTAGGCTTAGGCTTAACCTC

30

A C1/C2 short loop on chromosome 1 whose identifier is 134 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.2 and has the DNA sequence

TCTGCGTCTTTTCTCCCGCATTTTTTGTAGATCTACGTAGATCAAACCGAA
ATGAGGCACTTTCTGAATCCACGAGCTAGGCTTAAGCTTAGGCTTAAGCTT
AGGCCTTTTCTCAGGCTTAGGCTTAGGCTTA

5 A C1/C2 short loop on chromosome 1 whose identifier is 136 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.2 and has the DNA sequence

10 GCTTATGCTTGGGCTTAGGCTTAGGCGTAGGCTTAGGCTTAGGCTTAGGCT
TATGCTTAGACTTAGTCTCACTATCAGTCTTAGGCTTAGGCTTAGACTTAG
GCTTAAGCTTAGGCTTAAGCTTAGACTTAGGCTTAGGCTTAGGCTTAGGCT
TAGGCTTAGGTTTGGGCTTAGGCTTAGGCTTAACCTC

15 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

20 A C1/C2 short loop on chromosome 5 whose identifier is 21719 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene C39F7.5 and has the DNA sequence

ACGTTCTTAACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGC
ATTTTTGTAGATC

25 The match between the T1 sequence and the C1/C2 sequence is

ACGTTCTTAACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGC

The match between the T2 sequence and the C1/C2 sequence is

30

ACTCTGCGTCTCTTCTCCCGCATTTTTTGTAGATC

A C1/C2 short loop on chromosome 5 whose identifier is 21949 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene F16B4.4 and has the DNA sequence

5 ACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGCATTTTTTGT
AGATCTACGTAGATCAAGCCGAAATGAGACACTCTGACACCACG

The match between the T1 sequence and the C1/C2 sequence is

10 ACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGC

The match between the T2 sequence and the C1/C2 sequence is

15 ACTCTGCGTCTCTTCTCCCGCATTTTTTGTAGATC

A C1/C2 short loop on chromosome 5 whose identifier is 21655 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene C39F7.3 and has the DNA sequence

20 AACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGCATTTTTTG
TAGATCTACG

The match between the T1 sequence and the C1/C2 sequence is

25 AACCATGCAAAATCAGTTGAGAACTCTGCGTCTCTTCTCCCGC

The match between the T2 sequence and the C1/C2 sequence is


30 ACTCTGCGTCTCTTCTCCCGCATTTTTTGTAGATC

2. Many Connectrons control the expression of one set of genes in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Many different C1/C2 short loops can control the existence of one T1-T2 long loop. The C1/C2 short loops can be on the same chromosome or on different chromosomes from the T1-T2 long loop. This relationship is described as “many-to-one”. This relationship exists in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes

Example of a many-to-one connectron in prokaryotes – E. coli

In this example the existence of the T1-T2 (3197-3308) long loop is controlled by three C1/C2 short loops (3307, 3432 and 2218).



A double stranded DNA loop of length 93.542 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3197. This T1 control element has the DNA sequence

AAAAAATGCGCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3308. This T2 control element has the DNA sequence

5 TAAATTTCTCTTGTTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
ACACGGAACAACGGCAAACACGCCGCCGGGTCAGCGGGGTTCTCCTGAG
AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
GCGTATTATGCACACCCCGCGCCGCT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

10 rrsC gltU rrlC rrfC aspT trpT yifA yifE yifB
ilvL ilvG_1 ilvM ilvE ilvD ilvA ilvY ilvC ppiC
b3776 rep gppA rhlB trxA rhoL rho rfe wzzE
wecB rffH wecD wecE wzxE yifM_2 wecG yifK
argX hisR leuT proM aslB aslA hemY hemX
15 hemD cyaA cyaY b3808 dapF uvrD b3814 corA
yigF yigG rarD yigI pldA recQ yigJ yigK pldB
yigL yigM metR metE ysgA udp yigN ubiE yigP
b3836 yigU yigW_1 rfaH yigC ubiB fadA fadB
pepQ trkH hemG

20 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 1 whose identifier is 3307 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene hemG and has the DNA sequence

30 AAAAAATGCGCGGTCAGAAAATTATTTTAAATTTCTCTTGTTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAAGGCGTATTATG...GGAGTCTGCAAC
TCGACTCCATGAAGTCGGAATCGCTAGTAATCGTGGATCAGAATGCCACG

GTGAATACGTTCCCGGGCCTTGTACACACCGCCCGTCACACCATGGGAGT
GGGTTGCAAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCTTACCACT
TTGTGATTCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTAGGGGAAC
CTGCGGTTGGATCACCTCCTTACCTTAAAGAAGCGTTCTTTG

5

The match between the T1 sequence and the C1/C2 sequence is

AAAAAATGCGCGGTCAGAAAATTATTTTAAATTTCTCTTGTTCAGGCCGG
AATAACTCCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACG
CCGCCGGGTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAA
ATAAATGCTTGACTCTGTAGCGGGAA

10

The match between the T2 sequence and the C1/C2 sequence is

TAAATTTCTCTTGTTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
ACACGGAACAACGGCAAACACGCCGCCGGGTCAGCGGGGTTCTCCTGAG
AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
GCGTATTATGCACACCCCGCGCCGCT

15

A C1/C2 short loop on chromosome 1 whose identifier is 3432 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene *btuB* and has the DNA sequence

20

TGCGCGGTCAGAAAATTATTTTAAATTTCTCTTGTTCAGGCCGGAATAACT
CCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACGCCGCCGG
GTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAAATAAATG
CTTGACTCTGTAGCGGGAAGGCGTATTATGCACACC...ACACCATGGGAGT
GGGTTGCAAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCTTACCACT
TTGTGATTCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTAGGGGAAC
CTGCGGTTGGATCACCTCCTTACCTTAAAGAAGCGT

25

30

The match between the T1 sequence and the C1/C2 sequence is

5 TGC GCGGTCAGAAAATTATTTTAAATTTCTCTTGTCAGGCCGGAATAACT
 CCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACGCCGCCGG
 GTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAAATAAATG
 CTTGACTCTGTAGCGGGAA

The match between the T2 sequence and the C1/C2 sequence is

10 TAAATTTCTCTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTG
 ACACGGAACAACGGCAAACACGCCGCCGGGTCAGCGGGGTTCTCCTGAG
 AACTCCGGCAGAGAAAGCAAAAATAAATGCTTGACTCTGTAGCGGGAAG
 GCGTATTATGCACACCCCGCGCCGCT

15 A C1/C2 short loop on chromosome 1 whose identifier is 2218 controls the
 expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
 as a RNA single strand that is 3'UTR to the gene clpB and has the DNA sequence

20 CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
 AACGGCAAACACGCCGCCGGGC

The match between the T1 sequence and the C1/C2 sequence is


25 CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
 AACGGCAAACACGCCGCCGGGC

The match between the T2 sequence and the C1/C2 sequence is

30 CTTGTCAGGCCGGAATAACTCCCTATAATGCGCCACCACTGACACGGAAC
 AACGGCAAACACGCCGCCGGGC

Example of a many-to-one connectron in archea – *M. jannaschii*

In this example the existence of the T1-T2 (1630-1643) long loop is controlled by four C1/C2 short loops (1629, 1642, 124 and 1533).



A double stranded DNA loop of length 4.998 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1630. This T1 control element has the DNA sequence

TTATTAATTAGTTCAAAGGATTTTATTTAATTTCTAAGGGTTTGCTGGTTT
GATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTAAAAATTA
AGATTAATTAGGAAAGGAAATAAGATTTCTCTAACAGACAAGTTAAATTT
TTGGATTTAAAAAGATAAAAAT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1643. This T2 control element has the DNA sequence

TTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTATTG
AATTATTCAGATTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAAAT
TTCTCTAACAAATAAGTTAAATTTTGGATTTAAAAAGATAAAAATACTCT
GTTTTATTATGGAAAGAAAGAT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

MJ1597 MJ1598 MJ1599 MJ1600 MJ1601 MJ1602

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 1629 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1597 and has the DNA sequence

ATATGTTTGAAATTTGAAAATAAGAGTATTTAGAAGTTATTAATTAGTTCA
AAGGATTTTTATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATAT
TTGAGTTTATTGAATTATTCAGATTTTTTAAAAATTA

The match between the T1 sequence and the C1/C2 sequence is

TTATTAATTAGTTCAAAGGATTTTTATTTAATTTCTAAGGGTTTGCTGGTTT
GATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTTTAAAAATTA

The match between the T2 sequence and the C1/C2 sequence is

GCTGGTTTGATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTTA
AAAATTA

A C1/C2 short loop on chromosome 1 whose identifier is 1642 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1602 and has the DNA sequence

ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAA
ATTTCTCTAACAAATAAGTTAAATTTTTGGATTATAAAAAGATAAAAATACT
CTGTTTTATTATGGAAAGAAAGAT

The match between the T1 sequence and the C1/C2 sequence is

5 GCTGGTTTGATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTAA
AAAATTA

The match between the T2 sequence and the C1/C2 sequence is

10 TTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTATTG
AATTATTCAGATTTTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAAAT
TTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATAAAAATACTCT
GTTTTATTATGGAAAGAAAGAT

15 A C1/C2 short loop on chromosome 1 whose identifier is 124 controls the expression
of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA
single strand that is 3'UTR to the gene MJ0112 and has the DNA sequence

20 ATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTTTAAAAAT

The match between the T1 sequence and the C1/C2 sequence is

25 ATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTTTAAAAAT

The match between the T2 sequence and the C1/C2 sequence is

30 GCTGGTTTGATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTAA
AAAAT

A C1/C2 short loop on chromosome 1 whose identifier is 1533 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1486 and has the DNA sequence

5 TTTTTATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATATTTGAG
 TTTATT

The match between the T1 sequence and the C1/C2 sequence is

10 TTTTTATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATATTTGAG
 TTTATT

The match between the T2 sequence and the C1/C2 sequence is


15 GCTGGTTTGATTATTTAGAATATTTGAGTTTATT

Example of a many-to-one connectron in single-cell eukaryotes – *S. cerevisiae*

20 In this example the existence of the T1-T2 (5515-5533) long loop on chromosome 12
 is controlled by seventeen C1/C2 short loops (5516, 5532, 1939, 2323, 1942, 3286,
 3649, 4764, 4751, 5536, 6102, 8023, 7356, 3293, 3291, 3289 and 146).

- 25

5516 Chromosome 12
5532 Chromosome 12
1939 Chromosome 4
2323 Chromosome 5
1942 Chromosome 5
30
3286 Chromosome 7
3649 Chromosome 8
4764 Chromosome 12
4751 Chromosome 12
5536 Chromosome 13
35
6102 Chromosome 14



A double stranded DNA loop of length 6.466 kilo-bases on chromosome 12 is bounded on the left by a T1 sequence whose identifier is 5515. This T1 control element has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTIAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTG

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 5533. This T2 control element has the DNA sequence

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
 ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTIAGTGTTTGTTC
 ACGGCAGTAGCGAGAGACAAGTGGAAGAGTAGGATAAAAAGACAATC
 TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
 TCCGGGTAAGAGACAACAGGGCT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

YLR467W

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

5 A C1/C2 short loop on chromosome 12 whose identifier is 5516 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLR464W and has the DNA sequence

10 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

15 A C1/C2 short loop on chromosome 12 whose identifier is 5532 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLR467W and has the DNA sequence

20 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
25 AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

30 A C1/C2 short loop on chromosome 4 whose identifier is 1939 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed

as a RNA single strand that is 3'UTR to the gene YDR545W and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGG

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTATTTAGTGTGTTGTTGC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGG

A C1/C2 short loop on chromosome 5 whose identifier is 2323 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YER189W and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG

GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

5

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTGATTTAGTGTTTGTGTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

10

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTGTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTA
AGAGACAACAGGGCT

15

20

A C1/C2 short loop on chromosome 5 whose identifier is 1942 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YEL077C and has the DNA sequence

25

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTGATTTAGTGTTTGTGTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

30

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTGTCACGGCAGTAGCGAGAGACAAGTG
 5 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

10 ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTCTAGGGA
 ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTGTC
 ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
 TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
 TCCGGGTA
 15 AGAGACAACAGGGCT

A C1/C2 short loop on chromosome 7 whose identifier is 3286 controls the
 expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
 as a RNA single strand that is 3'UTR to the gene YGR296W and has the DNA
 20 sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTGTCACGGCAGTAGCGAGAGACAAGTG
 25 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

30 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTGTCACGGCAGTAGCGAGAGACAAGTG

GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

5

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

10

A C1/C2 short loop on chromosome 8 whose identifier is 3649 controls the
expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
as a RNA single strand that is 3'UTR to the gene YHR219W and has the DNA
sequence

15

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

20

The match between the T1 sequence and the C1/C2 sequence is

25

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

30

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

A C1/C2 short loop on chromosome 12 whose identifier is 4764 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLL066C and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

A C1/C2 short loop on chromosome 12 whose identifier is 4751 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLL067C and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTGTTGTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

A C1/C2 short loop on chromosome 13 whose identifier is 5536 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YML133C and has the DNA sequence

5 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

10 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

20 ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTCTAGGGA
 ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTC
 ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
 TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
 TCCGGGTAAGAGACAACAGGGCT

25 A C1/C2 short loop on chromosome 14 whose identifier is 6102 controls the
 expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
 as a RNA single strand that is 3'UTR to the gene YNL339C and has the DNA
 sequence

30 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGCGAGAGACAAGTG

GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is

5

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATGTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

10

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTGTGTTAGTGTGTTGTTGC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

15

A C1/C2 short loop on chromosome 16 whose identifier is 8023 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YPR204W and has the DNA sequence

20

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATGTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

25

30

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 5 AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
 10 ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTTAGTGTGTTG
 ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
 TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
 TCCGGGTAAGAGACAACAGGGCT

15 A C1/C2 short loop on chromosome 16 whose identifier is 7356 controls the
 expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
 as a RNA single strand that is 3'UTR to the gene YPL283C and has the DNA
 sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 20 ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 TTCACTGTTTTGATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

25 The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
 ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
 30 TTCACTGTTTTGATTTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
 GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
 AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTTGATTAGTGTTTGTTC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

A C1/C2 short loop on chromosome 8 whose identifier is 3293 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YHL050C and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTT

The match between the T1 sequence and the C1/C2 sequence is

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTT

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTT

A C1/C2 short loop on chromosome 8 whose identifier is 3291 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YHL050C and has the DNA sequence

ATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGC
GAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAA

The match between the T1 sequence and the C1/C2 sequence is

5

ATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGC
GAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAA

The match between the T2 sequence and the C1/C2 sequence is

10

ATGTAGTAGTATTTCACTGTTTTGATTTAGTGTTTGTTCACGGCAGTAGC
GAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAA

A C1/C2 short loop on chromosome 2 whose identifier is 145 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YBL113C and has the DNA sequence

15

CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAAC
ATCCGGGTAAGAGACAACAGGCT

20

The match between the T1 sequence and the C1/C2 sequence is

CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTG

25

The match between the T2 sequence and the C1/C2 sequence is

CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAAC
ATCCGGGTAAGAGACAACAGGCT

30

A C1/C2 short loop on chromosome 8 whose identifier is 3289 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed

as a RNA single strand that is 3'UTR to the gene YHL050C and has the DNA sequence

5 CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAAC
ATCCGGGTAAGAGACAACAGGCT

The match between the T1 sequence and the C1/C2 sequence is

10 CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

15 CTATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAAC
ATCCGGGTAAGAGACAACAGGCT

A C1/C2 short loop on chromosome 2 whose identifier is 146 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YBL113C and has the DNA sequence

20 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAA

The match between the T1 sequence and the C1/C2 sequence is


25 AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAA

The match between the T2 sequence and the C1/C2 sequence is

30 ATTATGTATTGTGTAGTATAGTATATTGTAAGAAA

Example of a many-to-one connectron in multi-cell eukaryotes – *C. elegans*

In this example the existence of the T1-T2 (3197-3308) long loop on chromosome 5 is controlled by three C1/C2 short loops (4382, 4375 and 28633).



A double stranded DNA loop of length 58.451 kilo-bases on chromosome 5 is bounded on the left by a T1 sequence whose identifier is 28632. This T1 control element has the DNA sequence

GCAAAAATTGACTGAAAATTTGAATTTCCCGCAAAAATTGACTGAAAAT
TTGAATTTCCCGCCAAAATTGACTGAAAATTTGAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 28697. This T2 control element has the DNA sequence

CAAAAATTGACTGAAAATTTGAATTTCCCTCCAAAATTGACTGAAAAT
TTGAATTTCCCGCCAAAATTGACTGAAAATTTGAATATCCCGCCAAAAA
TTGACTGAAAATTTGAATTTCCCGCCGAAAATTAAATGAAAATGGAATT
TCTCGCCGAA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

M162.8 M162.4 M162.3 M162.6 M162.2 M162.1 M162.7

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

5

A C1/C2 short loop on chromosome 1 whose identifier is 4382 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene Y43F8B.10 and has the DNA sequence

10

ATTATAGAAAATTTAAATTTCCCTCCAAAAAATTGACTGAAAATTTGAATT
TCCCTCCAAAAAATTGACTGAAAATTTGAATTTCCCGCCAAAAAATTGACTG
AAAATTTGAATATCCCGCCAAAAAATTGACTGAAAATTTGAATTTCCCGCC
GAAAATTAAATGAAAAATGGAATTTCTCGCCGAAAAATTCAGTAAAAATT
TGAATTTCTTGCCAAAAAATTGACTGAAAATTTGAATTTCTTGCCAAAAA
GTGACTGGGAATTTGAATTTCCCTCCAAAAAATTGACTGAAATTTTGAATTT
CCCGCTAAAAGTTGACT

15

The match between the T1 sequence and the C1/C2 sequence is

20

CAAAAATTGACTGAAAATTTGAATTTCCCGC

The match between the T2 sequence and the C1/C2 sequence is

25

CAAAAATTGACTGAAAATTTGAATTTCCCTCCAAAAAATTGACTGAAAAT
TTGAATTTCCCGCCAAAAAATTGACTGAAAATTTGAATATCCCGCCAAAAA
TTGACTGAAAATTTGAATTTCCCGCCGAAAAATTAAATGAAAAATGGAATT
TCTCGCCGAA

30

A C1/C2 short loop on chromosome 1 whose identifier is 4375 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed

as a RNA single strand that is 3'UTR to the gene Y43F8B.10 and has the DNA sequence

5 ATTATAGAAAATTTAAATTTCCCTCCAAAAAATTGACTGAAAATTTGAATT
TCCCTCCAAAAAATTGACTGAAAATTTGAATTTCCCGCCAAAAAATTGACTG
AAAATTTGAATATCCCGCCAAAAAATTGACTGAAAATTTGAATTTCCCGCC
GAAAATTAAATGAAAAATGGAATTTCTCGCCGAAAAATTCAGTAAAAATT
TGAATTTCTTGCCAAAAAATTGACTGAAAATTTGAATTTCTTGCCAAAAA
GTGACTGGGAATTTGAATTTCCCTCCAAAAAATTGACTGAAATTTTGAATTT
10 CCCGCTAAAAGTTGACT

The match between the T1 sequence and the C1/C2 sequence is

15 CAAAAATTGACTGAAAATTTGAATTTCCCGC

The match between the T2 sequence and the C1/C2 sequence is

20 CAAAAAATTGACTGAAAATTTGAATTTCCCTCCAAAAAATTGACTGAAAAT
TTGAATTTCCCGCCAAAAAATTGACTGAAAATTTGAATATCCCGCCAAAAA
TTGACTGAAAATTTGAATTTCCCGCCGAAAAATTAAATGAAAAATGGAATT
TCTCGCCGAA

25 A C1/C2 short loop on chromosome 5 whose identifier is 28633 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene M162.5 and has the DNA sequence

CAAAAATTGACTGAAAATTTGAATTTCCCGCAAAAAAATTGACTGAAAATT
TGAATTTCCCGCCAAAAAATTGACTGAAAATTTGAA

30 The match between the T1 sequence and the C1/C2 sequence is

CAAAAATTGACTGAAAATTTGAATTTCCCGCAAAAAATTGACTGAAAATT
TGAATTTCCCGCCAAAAATTGACTGAAAATTTGAA

The match between the T2 sequence and the C1/C2 sequence is

5

CAAAAATTGACTGAAAATTTGAATTTCCC

10


0966926.052004
100250" 52699360

3. One connectron controls the expression of many sets of genes in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

One C1/C2 short loop can control the existence of a many T1-T2 long loops. The C1/C2 short loop can be on the same chromosome or on different chromosomes from the T1-T2 long loops. This relationship is described as “one-to-many”. This relationship exists in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Example of a one-to-many connectron in prokaryotes – E. coli

In this example the existence of T1-T2 (3208-3315, 3436-3476, 3439-3478 and 3441-3479) long loops are controlled by one C1/C2 short loop (3206).



A double stranded DNA loop of length 93.377 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3208. This T1 control element has the DNA sequence

ACTCATCTTCGGGTGATGTTTGAGATATTTGCTCTTTAAAAATCTGGATCA
 AGCTGAAAATTGAAACACTGAACAACGAAAGTTGTTTCGTGAGTCTCTCAA
 ATTTTCGCAACACGATGATGAATCGAAAGAAACATCTTCGGGTTGTGAGG
 TTAAGCGACTAAGCGTACACGGTGGATGCCCTGGC...AGTGTGTTTCGACA
 CACTATCATTAAGTGAATCCATAGGTTAATGAGGCGAACCAGGGGGAAGTGA
 AAACATCTAAGTACCCCGAGGAAAAGAAATCAACCGAGATTCCCCCAGTA
 GCGGCGAGCGAACGGGGAGCAGCCCAGAGCCTGAATCAGT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3315. This T2 control element has the DNA sequence

TTTGCTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAACACTGAACAAC
 GAAAGTTGTTTCGTGAGTCTCTCAAATTTTCGCAACTCTGAAGTGAAACATC
 TTCGGGTTGTGAGGTTAAGCGACTAAGCGTACACGGTGGATGCCCTGGCA
 GTCAGAGGCGATGAAGGACGTGCTAATCTGCGATA...GGTTAATGAGGCG
 AACCAGGGGGAAGTGAACATCTAAGTACCCCGAGGAAAAGAAATCAACC
 GAGATTCCCCCAGTAGCGGCGAGCGAACGGGGAGCAGCCCAGAGCCTGA
 ATCAGTGTGTGTGTTAGTGGAAGCGTCTGGAAA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

rrlC	rrfC	aspT	trpT	yifA	yifE	yifB	ilvL	ilvG_1
ilvM	ilvE	ilvD	ilvA	ilvY	ilvC	ppiC	b3776	rep
gppA	rhlB	trxA	rhoL	rho	rfe	wzzE	wecB	rffH

	wecD	wecE	wzxE	yifM_2	wecG	yifK	argX	hisR
	leuT	proM	aslB	aslA	hemY	hemX	hemD	cyaA
	cyaY	b3808	dapF	uvrD	b3814	corA	yigF	yigG
	yigI	pIdA	recQ	yigJ	yigK	pIdB	yigL	yigM
5	metE	ysgA	udp	yigN	ubiE	yigP	b3836	yigU
	yigW_1	rfaH	yigC	ubiB	fadA	fadB	pepQ	trkH
	hemG	rrsA	ileT					

10 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 3206 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrsC and has the DNA sequence

15 GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTCTGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC
ACCTGCCTTAATATCTCAAACTCATCTTCGGGTGATGTTTGAGATATTTG
CTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAA...ACCGGCGATTTCCG
20 AATGGGGAAACCCAGTGTGTTTCGACACACTATCATTAAGTGAATCCATA
GGTTAATGAGGCGAACCGGGGGAAGTGAACATCTAAGTACCCCGAGGA
AAAGAAATCAACCGAGATTCCCCCAGTAGCGGCGAGCGAACGGGGAGCA
GCCAGAGCCTGAATCAGT

25 The match between the T1 sequence and the C1/C2 sequence is

ACTCATCTTCGGGTGATGTTTGAGATATTTGCTCTTTAAAAATCTGGATCA
AGCTGAAAATTGAAACACTGAACAACGAAAGTTGTTTCGTGAGTCTCTCAA
ATTTTCGCAACACGATGATGAATCGAAAGAAACATCTTCGGGTTGTGAGG
30 TTAAGCGACTAAGCGTACACGGTGGATGCCCTGGC...AGTGTGTTTCGACA
CACTATCATTAAGTGAATCCATAGGTTAATGAGGCGAACCGGGGGAAGT

AAACATCTAAGTACCCCGAGGAAAAGAAATCAACCGAGATTCCCCCAGTA
GCGGCGAGCGAACGGGGAGCAGCCCAGAGCCTGAATCAGT

The match between the T2 sequence and the C1/C2 sequence is

TTTGCTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAACACTGAACAAC
GAAAGTTGTTCTGTGAGTCTCTCAAATTTTCGCAAC

A double stranded DNA loop of length 41.279 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3436. This T1 control element has the DNA sequence

ACGCAACGCGTGATAAGCAATTTTCGTGTCCCCTTCGTCTAGAGGCCAG
GACACCGCCCTTTCACGGCGGTAACAGGGGTTCGAATCCCCTAGGGGACG
CCACTTGCTGGTT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3476. This T2 control element has the DNA sequence

AGTGAAAAGCAAGGCGTCTTGCGAAGCAGACTGATACGTCCCCTTCGTCT
AGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAGGGGTTCGAATCCC
CTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTCACCTGCCTTAATA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

gltT	rrlB	rrfB	murB	coaA	b3975	tyrU	thrT	tufB
secE	nusG	rplK	rplA	rplJ	rplL	rpoB	rpoC	htrC
thiH	thiF	thiE	yjaE	yjaD	hemE	nfi	yjaG	hupA
yjaH	yjaI	hydH	purD	purH				

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

5 A C1/C2 short loop on chromosome 1 whose identifier is 3206 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene *rrsC* and has the DNA sequence

10 GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTCGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC
ACCTGCCTTAATATCTCAAACTCATCTTCGGGTGATGTTTGAGATATTTG
CTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAACACTGAACAACGAAA
GTTGTTTCGTGAGTCTCTCAAATTTTCGCAACACGATGATGAATCGAAAGA
AACATCTTCGGGTTGTGAGGTTAAGCGACTAAGCGTACACGGTGGATGCC
15 CTGGCAGTCAGAGGCGATGAAGGACGTGCTAATCTGCGATAAGCGTCGGT
AAGGTGATATGAACCGTTATAACCGGCGATTTCGAATGGGGAAACCCAG
TGTGTTTCGACACACTATCATTAAGTGAATCCATAGGTTAATGAGGCGAA
CCGGGGGAACTGAAACATCTAAGTACCCCGAGGAAAAGAAATCAACCGA
GATTCCCCCAGTAGCGGCGAGCGAACGGGGAGCAGCCCAGAGCCTGAAT
20 CAGT

The match between the T1 sequence and the C1/C2 sequence is

25 GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTCGAATCCCCTAGGGGACGCCACTTGCTGGTT

The match between the T2 sequence and the C1/C2 sequence is

30 GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTCGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC
ACCTGCCTTAATA

5

A double stranded DNA loop of length 41.336 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3439. This T1 control element has the DNA sequence

CCTTAATATCTCAAAACTCATCTTCGGGTGATGTTTGAGATATTTGCTCTTT
AAAAATCTGGATCAAGCTGAAAATTGAAACACTGAACAACGA

10

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3478. This T2 control element has the DNA sequence

GTGATGTTTGAGATATTTGCTCTTTAAAAATCTGGATCAAGCTGAAAATTG
AAACACTGAACAACGAAAGTTGTCGTGAGTCTCTCAAATTTT

15

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

rrlB	rrfB	murB	coaA	b3975	tyrU	thrT	tufB	secE
nusG	rplK	rplA	rplJ	rplL	rpoB	rpoC	htrC	thiH
thiF	thiE	yjaE	yjaD	hemE	nfi	yjaG	hupA	yjaH
yjaI	hydH	purD	purH	gltV				

20

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25

A C1/C2 short loop on chromosome 1 whose identifier is 3206 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the generrsC and has the DNA sequence

30

GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTCGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC

ACCTGCCTTAATATCTCAAAACTCATCTTCGGGTGATGTTTGAGATATTTG
 CTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAA...ACCGGCGATTTCGG
 AATGGGGAAACCCAGTGTGTTTCGACACACTATCATTAAGTGAATCCATA
 GGTAAATGAGGCGAACCGGGGGAAGTGAACATCTAAGTACCCCGAGGA
 5 AAAGAAATCAACCGAGATTCCCCCAGTAGCGGCGAGCGAACGGGGAGCA
 GCCCAGAGCCTGAATCAGT

The match between the T1 sequence and the C1/C2 sequence is

10 CCTTAATATCTCAAAACTCATCTTCGGGTGATGTTTGAGATATTTGCTCTTT
 AAAAATCTGGATCAAGCTGAAAATTGAAACACTGAACAACGA

The match between the T2 sequence and the C1/C2 sequence is

15 GTGATGTTTGAGATATTTGCTCTTTAAAAATCTGGATCAAGCTGAAAATTG
 AAACACTGAACAACGAAAGTTGTTTCGTGAGTCTCTCAAATTTT

20 A double stranded DNA loop of length 38.285 kilo-bases on chromosome 1 is
 bounded on the left by a T1 sequence whose identifier is 3441. This T1 control
 element has the DNA sequence

25 AATTTTCGCAACACGATGATGAATCGAAAGAAACATCTTCGGGTTGTGAG
 GTTAAGCGACTAAGCGTACACGGTGGATGCCCTGGCAGTCAGAGGCGATG
 AAGGACGTGCTAATCTGCGATAAGCGTCGGTAAGGTGATATGAACCGTTA
 TAACCGGCGATTTCCGAATGGGGAAACCCAGTGTGT...GATGAGAGAAGA
 TTTTCAGCCTGATACAGATTAAATCAGAACGCAGAAGCGGTCTGATAAAA
 CAGAATTTGCCTGGCGGCAGTAGCGCGGTGGTCCCACCTGACCCCATGCC
 30 GAACTCAGAAGTGAAACGCCGTAGCGCCGATGGTAGTGTGGGGTCTCCCC
 ATGCGAG

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 3479. This T2 control element has the DNA sequence

5 AAGAAACATCTTCGGGTTGTGAGGTTAAGCGACTAAGCGTACACGGTGGATG
TGCCCTGGCAGTCAGAGGCGATGAAGGACGTGCTAATCTGCGATAAGCGT
CGGTAAGGTGATATGAACCGTTATAACCGGCGATTTCGAATGGGGAAAC
CCAGTGTGTTTCGACACACTATCATTAAGTGAATCC...CAGATTAAATCAG
AACGCAGAAGCGGTCTGATAAAACAGAATTTGCCTGGCGGCAGTAGCGC
GGTGGTCCCACCTGACCCCATGCCGAAGTCAGAAAGTGAAACGCCGTAGCG
10 CCGATGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGGAACTGCCAGGCA
TCAAATTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

15 rrlB rrfB murB coaA b3975 tyrU thrT tufB secE
nusG rplK rplA rplJ rplL rpoB rpoC htrC thiH
thiF thiE yjaE yjaD hemE nfi yjaG hupA yjaH
yjaI hydH purD purH gltV

20 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 1 whose identifier is 3206 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrsC and has the DNA sequence

30 GTCCCCTTCGTCTAGAGGCCAGGACACCGCCCTTTCACGGCGGTAACAG
GGGTTTCGAATCCCCTAGGGGACGCCACTTGCTGGTTTGTGAGTGAAAGTC
ACCTGCCTTAATATCTCAAACTCATCTTCGGGTGATGTTTGAGATATTTG
CTCTTTAAAAATCTGGATCAAGCTGAAAATTGAAA...ACCGGCGATTTCGG
AATGGGGAAACCCAGTGTGTTTCGACACACTATCATTAAGTGAATCCATA

GGTTAATGAGGCGAACCGGGGGAACCTGAAACATCTAAGTACCCCGAGGA
AAAGAAATCAACCGAGATTCCCCCAGTAGCGGCGAGCGAACCGGGGAGCA
GCCCAGAGCCTGAATCAGT

5 The match between the T1 sequence and the C1/C2 sequence is

AATTTTCGCAACACGATGATGAATCGAAAGAAACATCTTCGGGTTGTGAG
GTTAAGCGACTAAGCGTACACGGTGGATGCCCTGGCAGTCAGAGGCGATG
AAGGACGTGCTAATCTGCGATAAGCGTCGGTAAGGTGATATGAACCGTTA
10 TAACCGGCGATTTCGAATGGGGAAACCCAGTGTGTTTCGACACACTATC
ATTAAGTGAATCCATAGGTAAATGAGGCGAACCGGGGGAACCTGAAACATC
TAAGTACCCCGAGGAAAAGAAATCAACCGAGATTCCCCCAGTAGCGGCG
AGCGAACCGGGGAGCAGCCCAGAGCCTGAATCAGT

15 The match between the T2 sequence and the C1/C2 sequence is

AAGAAACATCTTCGGGTTGTGAGGTTAAGCGACTAAGCGTACACGGTGGG
TGCCCTGGCAGTCAGAGGCGATGAAGGACGTGCTAATCTGCGATAAGCGT
CGGTAAGGTGATATGAACCGTTATAACCGGCGATTTCGAATGGGGAAAC
20 CCAGTGTGTTTCGACACACTATCATTAAGTGAATCCATAGGTAAATGAGGC
GAACCGGGGGAACCTGAAACATCTAAGTACCCCGAGGAAAAGAAATCAAC
CGAGATTCCCCCAGTAGCGGCGAGCGAACCGGGGAGCAGCCCAGAGCCTG
AATCAGT


25 -----

Example of a one-to-many connectron in archea – *M. jannaschii*


30 In this example the existence of T1-T2 (534-611, 1139-1159, and 1630-1643) long
loops are controlled by one C1/C2 short loop (1642).

1642 Chromosome 1


5



10



15



20

A double stranded DNA loop of length 72.886 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 534. This T1 control element has the DNA sequence

25

TAAGTAAATAAAATTTCTCTAACAAATAAGTTAAATT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 611. This T2 control element has the DNA sequence

30

TAAATAAAATTTCTCTAACAAATAAGTTAAATTTTGGATTAAAAAGATA
AAAATGCT

35

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

MJ0486	MJ0487	MJ0488	MJ0489	MJ0490	MJ0492	MJ0493
MJ0494	MJ0495	MJ0496	MJ0497	MJ0499	MJ0500	MJ0501

MJ0502	MJ0503	MJ0504	MJ0506	MJ0507	MJ0508	MJ0509
MJ0510	MJ0511	MJ0512	MJ0513	MJ0514	MJ0514	MJ0517
MJ0519	MJ0520	MJ0521	MJ0522	MJ0523	MJ0525	MJ0526
MJ0526	MJ0529	MJ0530	MJ0531	MJ0532	MJ0534	MJ0535
MJ0536	MJ0538	MJ0539	MJ0540	MJ0541	MJ0542	MJ0543
MJ0544	MJ0545	MJ0547	MJ0548	MJ0549	MJ0550	MJ0552
MJ0553	MJ0554	MJ0555	MJ0556	MJ0558	MJ0559	MJ0560
MJ0561	MJ0562	MJ0563	MJ0564			

10 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

15 A C1/C2 short loop on chromosome 1 whose identifier is 1642 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1602 and has the DNA sequence

ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAA
ATTTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATAAAAAATACT
20 CTGTTTTATTATGGAAAGAAAGAT

The match between the T1 sequence and the C1/C2 sequence is

AAGTAAATAAAATTTCTCTAACAAATAAGTTAAATT

25

The match between the T2 sequence and the C1/C2 sequence is

TAAATAAAATTTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATA
AAAAT

30

A double stranded DNA loop of length 14.509 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1139. This T1 control element has the DNA sequence

5 ATTTATTAATTAGTTCAAAGGATTTTTATTTAATTTCTAAGGGTTAGCTGG
 TTTGATTGTTTAAAATATTTGAGTTTA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1159. This T2 control element has the DNA sequence

10 ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
 TGAATTATTCAGATTTTAAAAAATTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

15 MJ1096 MJ1097 tRNA-Arg-3 MJ1098 MJ1099 MJ1100 MJ1101
 MJ1102 MJ1103 MJ1104 MJ1105 MJ1106 MJ1107 MJ1108

20 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 1 whose identifier is 1642 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1602 and has the DNA sequence

30 ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
 TGAATTATTCAGATTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAA
 ATTTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATAAAAATACT
 CTGTTTTATTATGGAAAGAAAGAT

The match between the T1 sequence and the C1/C2 sequence is

ATTTAATTTCTAAGGGTTAGCTGGTTTGATT

The match between the T2 sequence and the C1/C2 sequence is

ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTAAAAATTA

A double stranded DNA loop of length 4.998 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1630. This T1 control element has the DNA sequence

TTATTAATTAGTTCAAAGGATTTTATTTAATTTCTAAGGGTTTGCTGGTTT
GATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTAAAAATTA
AGATTAATTAGGAAAGGAAATAAGATTTCTCTAACAGACAAGTTAAATTT
TTGGATTTAAAAAGATAAAAAT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1643. This T2 control element has the DNA sequence

TTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTATTG
AATTATTCAGATTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAAAT
TTCTCTAACAAATAAGTTAAATTTTGGATTTAAAAAGATAAAAATACTCT
GTTTTATTATGGAAAGAAAGAT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

MJ1597 MJ1598 MJ1599 MJ1600 MJ1601 MJ1602

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 1642 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1602 and has the DNA sequence

ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAA
ATTTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATAAAAAATACT
CTGTTTTATTATGGAAAGAAAGAT

The match between the T1 sequence and the C1/C2 sequence is

GCTGGTTTGATTATTTAGAATATTTGAGTTTATTGAATTATTCAGATTTTAA
AAAATTA


The match between the T2 sequence and the C1/C2 sequence is

TTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTATTG
AATTATTCAGATTTTTAAAAATTAGGATTAATTAGGCAAGTAAATAAAAT
TTCTCTAACAAATAAGTTAAATTTTTGGATTAAAAAGATAAAAAATACTCT
GTTTTATTATGGAAAGAAAGAT

Example of a one-to-many connectron in single-cell eukaryotes – *S. cerevisiae*

In this example the existence of T1-T2 (158-171, 293-317, 4295-4308 and 5916-5923) long loops are controlled by one C1/C2 short loop (86).

86 Chromosome 1



A double stranded DNA loop of length 20.391 kilo-bases on chromosome 2 is bounded on the left by a T1 sequence whose identifier is 158. This T1 control element has the DNA sequence

CCAATTGTTGGAATAAAAATCAACTATCATCTACTAACTAGTATTTACGTT
ACTAGTATATTATCATATACGGTGTTAGAAGATGACGCAAATGATGAGAA
ATAGTCATCTAAATTAGTGGAAGCTGAAACGCAAGGATTGATAATGTAAT
AG

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 171. This T2 control element has the DNA sequence

ATAATTGTTGGAATAAAAATCAACTATCATCTACTAACTAGTATTTACGTT
 ACTAGTATATTATCATATACGGTGTTAGAAGATGACACAAATGATGAGAA
 ATAGTCATCTAAATTAGTGGAAGCTGAAACGCAAGGATTGATAATGTAAT
 5 AGGATCAATGAATATTAACATATAAAAATGATGATAATAATA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

10 YBL107W-A TL(UAA)B1 YBL107C YBL106C YBL105C YBL104C
 YBL103C YBL102W YBL101C

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

15

A C1/C2 short loop on chromosome 1 whose identifier is 86 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YAR009C and has the DNA sequence

20 ATCTATTACATTATGGGTGGTATGTTGGAATAGAAATCAACTATCATCTAC
 TAACTAGTATTTACATTACTAGTATATTATCATATACGGTGTTAGAAGATG
 ACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAAACGCA
 AGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGAAT
 GAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGG
 25 ATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATATT
 ATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCACCCAT
 TTCTCAGAA

The match between the T1 sequence and the C1/C2 sequence is

30

AAATCAACTATCATCTACTAACTAGTATTTAC

The match between the T2 sequence and the C1/C2 sequence is

AAATCAACTATCATCTACTAACTAGTATTTAC

5

A double stranded DNA loop of length 38.470 kilo-bases on chromosome 2 is bounded on the left by a T1 sequence whose identifier is 293. This T1 control element has the DNA sequence

10

GAATTGTTGGAATAAAAATCCACTATCGTCTATCAACTAATAGTTATATTA
TCAATATATTATCATATACGGTGTTAAGATGATGACATAAGTTATGAGAA
GCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCAAGGATTGATAATGTAAT
AGGATAATGAAACATATAAAACGGAATGAGGAATAATCGTAATATTAGT
ATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTTGAGGAGAAC
TTCTAGT

15

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 317. This T2 control element has the DNA sequence

20

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTC
GAGGAGAACTTCTAGTATATTCTGTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

25

YBL005W-B TS(AGA)B YBL004W YBL003C YBL002W YBL001C
YBR001C YBR002C YBR003W YBR004C YBR005W YBR006W
YBR007C YBR008C YBR009C YBR010W YBR011C YBR012C

30

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 86 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YAR009C and has the DNA sequence

5

ATCTATTACATTATGGGTGGTATGTTGGAATAGAAATCAACTATCATCTAC
TAACTAGTATTTACATTACTAGTATATTATCATATACGGTGTTAGAAGATG
ACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAAACGCA
AGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGAAT
10 GAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGG
ATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATATT
ATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCACCCAT
TTCTCAGAA

15

The match between the T1 sequence and the C1/C2 sequence is

AAACATATAAAACGGAATGAGGAATAATCGTAATATTAGTATGTAGAAAT
ATAGATTCCATTTTGAGGATTCCTATATCCT

20

The match between the T2 sequence and the C1/C2 sequence is

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTC
GAGGAGAACTTCTAGTATATTCTGTA

25

A double stranded DNA loop of length 11.020 kilo-bases on chromosome 10 is bounded on the left by a T1 sequence whose identifier is 4295. This T1 control element has the DNA sequence

30

AAACGCAAGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAA
ACGGAATGAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCAT
TTTGAGGATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTG

5 This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 4308. This T2 control element has the DNA sequence

GGAAGCTGAAACGCAAGGATTGATAATGTAATAGGATCAATGAATATAA
ACATATAAAACGGAATGAGGAATAATCGTAATATTAGTATGTAGAAATAT
10 AGATTCCATTTTGAGGATTCCTATATCCTCGAGGAGAACTTCTAGTATATT
CTGTATACCTAATATTATAGCCTTTATCAA

This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

15 YJR027W YJR029W

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2
short loops.

20 A C1/C2 short loop on chromosome 1 whose identifier is 87 controls the expression
of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA
single strand that is 3'UTR to the gene YAR009C and has the DNA sequence

25 ATCTATTACATTATGGGTGGTATGTTGGAATAGAAATCAACTATCATCTAC
TAACTAGTATTTACATTACTAGTATATTATCATATACGGTGTTAGAAGATG
ACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAAACGCA
AGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGAAT
GAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGG
30 ATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATATT
ATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCACCCAT
TTCTCA

A double stranded DNA loop of length 5.462 kilo-bases on chromosome 13 is bounded on the left by a T1 sequence whose identifier is 5916. This T1 control element has the DNA sequence

AAGCTGAAGTGCAAGGATTGATAATGTAATAGGATAATGAAACATATAA
AACGGAATGAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCA
TTTTGAGGATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 5923. This T2 control element has the DNA sequence

TAATAGGATAATGAAACATATAAAACGGAATGAGGAATAATCGTAATATT
AGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTCGAGGAG
AACTTCTAGTATATTCTGTATACCTAATATTATAGCCTTTATCAA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

YML045W

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 87 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YAR009C and has the DNA sequence



ATCTATTACATTATGGGTGGTATGTTGGAATAGAAATCAACTATCATCTAC
TAACTAGTATTTACATTACTAGTATATTATCATATACGGTGTTAGAAGATG

ACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAAACGCA
 AGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGAAT
 GAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGG
 ATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATATT
 5 ATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCACCCAT
 TTCTCA

10 Example of a one-to-many connectron in multi-cell eukaryotes – *C. elegans*

In this example the existence of T1-T2 (16554-16661 and 21565-21590) long loops are controlled by one C1/C2 short loop (21591).

15



30

A double stranded DNA loop of length 50.159 kilo-bases on chromosome 4 is
 35 bounded on the left by a T1 sequence whose identifier is 16554. This T1 control
 element has the DNA sequence

TGCCTGAAAAAATTGGCTCCGAGTTAGGACACTTGGGGTGGTCAAAAAAT
TTTGTGACTATTGTCAAATGAAAGATCATAGTTGATAACATAAATTCCCAA
AGTTTCATAAAAAATCGATACGCAGCGAACAAAGTTATCAATT

- 5 This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 16661. This T2 control element has the DNA sequence

CACTTGGGGTGGTCAAAAAATTTTGTGATTATTGTCAAATGAAAGATCAT
GGTTGATAACATAAATTCCCAAAGTTTCATAAAAAATCGATACGCAGCGAA
10 CAAAGTTATGATTTTTGACCCGGAACCTTATTTGGAGACCTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

15 C23H5.7 C23H5.8a C23H5.3 C23H5.2 C23H5.9 C23H5.1

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

- 20 A C1/C2 short loop on chromosome 5 whose identifier is 21591 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene F25A2.1 and has the DNA sequence

TATTGTCAAATGAAAGATCATGGTTGATAACATAAATTCCCACAATTTTCAT
25 AAAAAATCGATACGCAGCGAACAAAGTTATGATTTTTGACCCGGAACCTTAT
TTGGAGACCTAATATT

The match between the T1 sequence and the C1/C2 sequence is

30 TTTCATAAAAAATCGATACGCAGCGAACAAAGTTAT

The match between the T2 sequence and the C1/C2 sequence is

TATTGTCAAATGAAAGATCATGGTTGATAACATAAAATTCCCA

5

A double stranded DNA loop of length 18.142 kilo-bases on chromosome 5 is bounded on the left by a T1 sequence whose identifier is 21565. This T1 control element has the DNA sequence

10

CTCCGAGTTAGGACACTTGGGGTGGACAAAAAATTTTGTGACTATTGTCA
AATGAAAGATCATGGTTGATAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 21590. This T2 control element has the DNA sequence

15

TATTGTCAAATGAAAGATCATGGTTGATAACATAAAATTCCCACAATTCAT
AAAAATCGATACGCAGCGAACAAGTTATGATTTTTGACCCGGAACCTTAT
TTGGAGACCTAATA

20

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

T21H3.2 T21H3.1 F25A2.1

25

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 5 whose identifier is 21591 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene F25A2.1 and has the DNA sequence

30

096696-05001

TATTGTCAAATGAAAGATCATGGTTGATAACATAAATTCCCACAATTCAT
AAAAATCGATACGCAGCGAACAAGTTATGATTTTTGACCCGGAAGTTAT
TTGGAGACCTAATATT

5 The match between the T1 sequence and the C1/C2 sequence is

TATTGTCAAATGAAAGATCATGGTTGATAA

The match between the T2 sequence and the C1/C2 sequence is

10

TATTGTCAAATGAAAGATCATGGTTGATAACATAAATTCCCACAATTCAT
AAAAATCGATACGCAGCGAACAAGTTATGATTTTTGACCCGGAAGTTAT
TTGGAGACCTAATA

15


096695-0530
100250-6269860

4. Connectrons occur between prokaryotes and their plasmids.

Connectron relationships exist between prokaryotes and their plasmids. These connectrons implement a control mechanism between the two genomes that makes it possible for them to form a symbiotic relationship. In the case of *D. radiodurans* the relationship is not symmetric. The *D. radiodurans* genome sends C1/C2 short loops to the MP1 plasmid.

Example of a prokaryote/plasmid connectron – *D. radiodurans*

In this example the existence of T1-T2 (2654-2694 and 2692-2749) long loops in chromosome 3 that is the plasmid MP1 are controlled by one C1/C2 short loop (16) in chromosome 1.



A double stranded DNA loop of length 46.903 kilo-bases on chromosome 3 (plasmid MP1) is bounded on the left by a T1 sequence whose identifier is 2654. This T1 control element has the DNA sequence

CAGCGTTTTTCTCGCTGTTCTGACGGCTGAACGCCCTGAATCTCTCCCG
 GTATGCAGCCTGCTCGGAGAGTACGATTCTGTCGTTGGCTGCACCGAAGTG
 ACGATGGGGCCATTCCGTGGGGCGCGTTACACCAGGCGACTGTCAGTACA
 5 GCAATCGAGAGTGGGCTGATCAGCCCACTGTGCGTTCTGGCCATCGACGC
 CTCTTTTCACCGCAAAGCCGGTCAGCACACCGCACACCTCGGCTCGTTCTG
 GAATGGCTGTGCCGCGCGGACC

10 This double stranded DNA loop is bounded on the right by a T2 control element
 whose identifier is 2694. This T2 control element has the DNA sequence

GCTGAACGCCCTGAATCTCTCCCGGTATGCAGCCTGCTCGGAGAGTACGA
 TTCGTCGTTGGCTGCACCGAAGTGACGATGGGGCCATTCCGTGGGGCGCG
 TTACACCAGGCGACTGTCAGTACAGCAATCGAGAGTGGGCTGATCAGCCC
 15 ACTGTGCGTTCTGGCCATCGACGCCTCTTTTCACCGCAAAGCCGGTCAGCA
 CACCGCACACCTCGGCTCGTTCTGGAATGGCTGTGCCGCGCGGACCGAAC
 GCGGAATCGAGCAATCCTGTTGT

20 This long T1/T2 double stranded DNA loop modulates the expression of the
 following genes

DRB0020	DRB0021	DRB0022	DRB0023	DRB0024	DRB0025
DRB0027	DRB0030	DRB0032	DRB0033	DRB0034	DRB0035
DRB0037	DRB0038	DRB0039	DRB0041	DRB0042	DRB0043
25 DRB0044	DRB0045	DRB0047	DRB0051	DRB0052	DRB0054
DRB0055	DRB0057				

30 This long T1/T2 double stranded DNA loop modulates the expression of the
 following C1/C2 short loops

A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2693
 controls the expression of the genes of one or more other T1/T2 long loops. This

C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DRB0057 and has the DNA sequence

5 CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTTTCCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTTCCTGG
AC

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

10

A C1/C2 short loop on chromosome 1 whose identifier is 16 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DR0009 and has the DNA sequence

15 GCTGTGAAATCACCGCTTCCAATGGGTCTGATGGCCATCCTACAGTACGTT
CTCAGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTTCCTGACCGTGCT
GCTCAGCGTTTTTCTCGCTGTTTCCTGGACGGCTGAACGCCCTGAATCTCTC
CCGGTATGCAGCCTGCTCGGAGAGTACGATTTCGT

20

The match between the T1 sequence and the C1/C2 sequence is

25 CAGCGTTTTTCTCGCTGTTTCCTGGACGGCTGAACGCCCTGAATCTCTCCCG
GTATGCAGCCTGCTCGGAGAGTACGATTTCGTCGTTGGCTGCACCGAAGTG
ACGATGGGGCCATTCCGTGGGGCGCGTTACACCAGGCGACTGTCAGTACA
GCAATCGAGAGTGGGCTGATCAGCCCACTGTGCGTTCTGGCCATCGACGC
CTCTTTTCACCGCAAAGCCGGTCAGCACACCGCACACCTCGGCTCGTTCTG
GAATGGCTGTGCCGCGCGGACC

30

The match between the T2 sequence and the C1/C2 sequence is

5 GCTGAACGCCCTGAATCTCTCCCGGTATGCAGCCTGCTCGGAGAGTACGA
 TTCGTCGTTGGCTGCACCGAAGTGACGATGGGGCCATTCCGTGGGGCGCG
 TTACACCAGGCGACTGTCAGTACAGCAATCGAGAGTGGGCTGATCAGCCC
 ACTGTGCGTTCTGGCCATCGACGCCTCTTTTCACCGCAAAGCCGGTCAGCA
 CACCGCACACCTCGGCTCGTTCTGGAATGGCTGTGCCGCGCGGACCGAAC
 GCGGAATCGAGCAATCCTGTTGT

10 A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2768
 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is
 expressed as a RNA single strand that is 3'UTR to the gene DRB0133 and has the
 DNA sequence

15 GCTGTGAAATCACCGCTTCCAATGGGTCTGATGGCCATCCTACAGTACGTT
 CTCAGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTCTGACCGTGCT
 GCTCAGCGTTTTTCTCGCTGTTCTGGACGGCTGAACGCCCTGAATCTCTC
 CCGGTATGCAGCCTGCTCGGAGAGTACGATTCTGT

20 The match between the T1 sequence and the C1/C2 sequence is

25 CAGCGTTTTTCTCGCTGTTCTGGACGGCTGAACGCCCTGAATCTCTCCCG
 GTATGCAGCCTGCTCGGAGAGTACGATTCTGTCGTTGGCTGCACCGAAGTG
 ACGATGGGGCCATTCCGTGGGGCGCGTTACACCAGGCGACTGTCAGTACA
 GCAATCGAGAGTGGGCTGATCAGCCCACTGTGCGTTCTGGCCATCGACGC
 CTCTTTTCACCGCAAAGCCGGTCAGCACACCGCACACCTCGGCTCGTTCTG
 GAATGGCTGTGCCGCGCGGACC

The match between the T2 sequence and the C1/C2 sequence is

30 GCTGAACGCCCTGAATCTCTCCCGGTATGCAGCCTGCTCGGAGAGTACGA
 TTCGTCGTTGGCTGCACCGAAGTGACGATGGGGCCATTCCGTGGGGCGCG
 TTACACCAGGCGACTGTCAGTACAGCAATCGAGAGTGGGCTGATCAGCCC

A double stranded DNA loop of length 68.612 kilo-bases on chromosome 3 (plasmid MP1) is bounded on the left by a T1 sequence whose identifier is 2692. This T1 control element has the DNA sequence

CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTTCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTCTGG
AC

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 2749. This T2 control element has the DNA sequence

AGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTCTGACCGTGCTGCT
CAGCGTTTTTCTCGCTGTTCTCTGGACGGCTGAACGCCCTGAATCTCTCCCG
GT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

DRB0059	DRB0060	DRB0061	DRB0062	DRB0064	DRB0065
DRB0066	DRB0067	DRB0068	DRB0069	DRB0070	DRB0072
DRB0073	DRB0074	DRB0076	DRB0077	DRB0079	DRB0080
DRB0081	DRB0083	DRB0085	DRB0086	DRB0087	DRB0088
DRB0089	DRB0090	DRB0092	DRB0093	DRB0094	DRB0096
DRB0097	DRB0098	DRB0102	DRB0103	DRB0104	DRB0105
DRB0106	DRB0107	DRB0111	DRB0112		

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2693 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DRB0057 and has the DNA sequence

5

CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTTCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTGG
AC

10

A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2695 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DRB0057 and has the DNA sequence

15

GCTGAACGCCCTGAATCTCTCCCGGTATGCAGCCTGCTCGGAGAGTACGA
TTCGTCGTTGGCTGCACCGAAGTGACGATGGGGCCATTCCGTGGGGCGCG
TTACACCAGGCGACTGTCAGTACAGCAATCGAGAGTGGGCTGATCAGCCC
ACTGTGCGTTCTGGCCATCGACGCCTCTTTTCACCGCAAAGCCGGTCAGCA
CACCGCACACCTCGGCTCGTTCTGGAATGGCTGTGCCGCGCGGACCGAAC
20 GCGGAATCGAGCAATCCTGTTGT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25

A C1/C2 short loop on chromosome 1 whose identifier is 16 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DR0009 and has the DNA sequence

30

GCTGTGAAATCACCGCTTCCAATGGGTCTGATGGCCATCCTACAGTACGTT
CTCAGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTCTGACCGTGCT
GCTCAGCGTTTTTCTCGCTGTTCTGGACGGCTGAACGCCCTGAATCTCTC
CCGGTATGCAGCCTGCTCGGAGAGTACGATTCGT

The match between the T1 sequence and the C1/C2 sequence is

5 CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTTCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTCTGG
AC

The match between the T2 sequence and the C1/C2 sequence is

10 AGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTCTGACCGTGCTGCT
CAGCGTTTTTCTCGCTGTTCTGACGGCTGAACGCCCTGAATCTCTCCCG
GT

15 A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2768
controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene DRB0133 and has the
DNA sequence

20 GCTGTGAAATCACCGCTTCCAATGGGTCTGATGGCCATCCTACAGTACGTT
CTCAGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTTCTGACCGTGCT
GCTCAGCGTTTTTCTCGCTGTTCTGACGGCTGAACGCCCTGAATCTCTC
CCGGTATGCAGCCTGCTCGGAGAGTACGATTCGT...CGGACCGAACGCGGA
ATCGAGCAATCCTGTTGTGCCCTCATTGATGTCCAGCACCGGCAGGCCTTG
25 ACGGTCGATGTCCGTCAGACCCTGACCGGGTCTGAGGCTCCAACTCGTCT
GGAACAG

The match between the T1 sequence and the C1/C2 sequence is

30 CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTTCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTCTGG
AC

The match between the T2 sequence and the C1/C2 sequence is

AGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTCCTGACCGTGCTGCT
CAGCGTTTTTCTCGCTGTTCTGACGGCTGAACGCCCTGAATCTCTCCCG
GT

A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2693 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DRB0057 and has the DNA sequence

CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTCCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTG
AC

The match between the T1 sequence and the C1/C2 sequence is

CTGATGGCCATCCTACAGTACGTTCTCAGCGCGGTCCCGCTGCGCAAGAC
GCAGCGGAATTCCTGACCGTGCTGCTCAGCGTTTTTCTCGCTGTTCTG
AC

The match between the T2 sequence and the C1/C2 sequence is

AGCGCGGTCCCGCTGCGCAAGACGCAGCGGAATTCCTGACCGTGCTGCT
CAGCGTTTTTCTCGCTGTTCTGAC


A C1/C2 short loop on chromosome 3 (plasmid MP1) whose identifier is 2653 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene DRB0017 and has the DNA sequence

5. Connectrons occur in plants and higher animals

Connectron relationships exist in plant and higher animals.

5 Example of a plant connectron – *A. thaliana*

In this example the existence of the T1-T2 (423-469) long loop is controlled by six C1/C2 short loops (972, 21396, 422, 21762, 21813 and 10882). The T1-T2 long loop controls the expression of six genes on chromosome 2 in addition to two C1/C2 (426 and 430) short loops.



A double stranded DNA loop of length 42.285 kilo-bases on chromosome 2 is bounded on the left by a T1 sequence whose identifier is 423. This T1 control element has the DNA sequence

TATCTCTTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAATTTA
ATTAAAAAACGAAATA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 469. This T2 control element has the DNA sequence

TACTAATTTAATTAATTAATAATTTAATTAAAAACGAAATACATTATTAATT
TTCAAAAATAATAACC

5 This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

At2g02070 At2g02080 At2g02090 At2g02100 At2g02120 At2g02130

10 This long T1/T2 double stranded DNA loop modulates the expression of the
following C1/C2 short loops

15 A C1/C2 short loop on chromosome 2 whose identifier is 426 controls the expression
of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene At2g02060 and has the
DNA sequence

TTCCAAAATAATAACCAATCAAAATCAACATATAAGATTTGATATCTAA
ATTTT

20 A C1/C2 short loop on chromosome 2 whose identifier is 430 controls the expression
of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene At2g02060 and has the
DNA sequence

25 TTGCGGAAAAATAATATCATCATTATAAAAAATAATTAGAGTTTTTCGC
ATAT

30 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2
short loops.

[illegible]

5

10

10

15

15

20

25

30

30

The match between the T2 sequence and the C1/C2 sequence is

TACTAATTTAATTAATTAAATTTAATTAAAAACGAAATACATTATTAATT
T

5

A C1/C2 short loop on chromosome 2 whose identifier is 422 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene At2g02060 and has the DNA sequence

10

TAACCTTAATTTTTGTAAGTAATTATATAGGTATGCCATTAGAAATAAAAT
TTTAAAGAGTAAATTAATTTATCTCTTTAAGGATTAAAAAGTCAAATACTA
ATTTAATTAATTAATTTAATTAATAAAAAACGAAATA

The match between the T1 sequence and the C1/C2 sequence is

15

TATCTCTTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTA
ATTAAAAAACGAAATA

The match between the T2 sequence and the C1/C2 sequence is

20

TACTAATTTAATTAATTAAATTTAATTAAAAACGAAATA

A C1/C2 short loop on chromosome 4 whose identifier is 21762 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene AT4g17510 and has the DNA sequence

25

TTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTAATTAAA
AACGAAATACATT

30

The match between the T1 sequence and the C1/C2 sequence is

TTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTAATTAAA
AAACGAAATA

The match between the T2 sequence and the C1/C2 sequence is

TACTAATTTAATTAATTAAATTTAATTAAAAACGAAATACATT

A C1/C2 short loop on chromosome 4 whose identifier is 21813 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene AT4g17680 and has the DNA sequence

TTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTAATTAAA
AAACGAAATACATT

The match between the T1 sequence and the C1/C2 sequence is

TTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTAATTAAA
AAACGAAATA

The match between the T2 sequence and the C1/C2 sequence is

TACTAATTTAATTAATTAAATTTAATTAAAAACGAAATACATT

A C1/C2 short loop on chromosome 2 whose identifier is 10882 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene At2g26540 and has the DNA sequence

TATCTCTTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTA
ATTAA

The match between the T1 sequence and the C1/C2 sequence is

TATCTCTTTAAGGATTAAAAAGTCAAATACTAATTTAATTAATTAAATTTA
ATTAA

5

The match between the T2 sequence and the C1/C2 sequence is

TACTAATTTAATTAATTAAATTTAATTAA

10

Example of a animal connectron – D. megalomaster

A double stranded DNA loop of length 88.159 kilo-bases on chromosome 4 is
bounded on the left by a T1 sequence whose identifier is 3340. This T1 control
element has the DNA sequence

ACCTAAAAGAAGTACCGTTTTTTACTCCTAATTACCAATTCTAACCATCCA
TATCACTTTTTGACGGACTCCGTGAAAATAATTTTTGGCCAAATTTTCGCA
TTTTTTGTAAGGGGTAACATCATAAAAATT

This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 3372. This T2 control element has the DNA sequence

AAAAAAGTACCGCGTTTTTTACTCCTAATTACCAATTCTAACCATCCATATCA
CTTTTTGACGGACTCCGTGAAAATAATTTTTGGCCAAATTTTCGCATTTTTT
GTAAGGGGTAACATCATCAAAATTTGCGAAAAA

This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

[Some of the following gene names have not been determined.]

-	-	-	-	-
-	CG11207	-	CG2186	CG2157
-	Ork1	-	-	-
-				

5

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

10 A C1/C2 short loop on chromosome 4 whose identifier is 3362 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene XXX and has the DNA sequence

15 AAAAAAGTACCGCGTTTTACTCCTAATTACCAATTCTAACCATCCATATCA
CTTTTTGACGGACTCCGTTAAAATAATTTTTGACCAAATTTTCGCATTTTTT
GTAATCAAAATTTGCAAAAAATTGAAAAAAC

20 A C1/C2 short loop on chromosome 4 whose identifier is 3364 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene XXX and has the DNA sequence

25 CAAAATTTGAATGCAAATCGATTGGGAATCAAAAAACAAACTCAACGAG
GTATGACATTCCATATTTGGGCCATTATTTCCAA

30 A C1/C2 short loop on chromosome 4 whose identifier is 3366 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene XXX and has the DNA sequence

TTTTTTCACAAAAATTAGGAAAATGATTTTGGGTAAAAAATGAATATTT
AAGTTGGGTTTT

A C1/C2 short loop on chromosome 4 whose identifier is 3369 controls the
expression of the genes of one or more other T1/T2 long loops. This C1/C2 short
loop is expressed as a RNA single strand that is 3'UTR to the gene XXX and has the
DNA sequence

AAATCGATTGGGAATCAAAAAACAAACCTCAACGAGGTATGACATTCCAT
ATCTGGGCCATTATTTCCAATCTTTTGATCAAAATAC

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2
short loops.

A C1/C2 short loop on chromosome 4 whose identifier is 3373 controls the
expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
as a RNA single strand that is 3'UTR to the gene XXX and has the DNA sequence

AAAAAAGTACCGCGTTTTACTCCTAATTACCAATTCTAACCATCCATATCA
CTTTTTGACGGACTCCGTGAAAATAATTTTTGGCCAAATTTTCGCATTTTTT
GTAAGGGGTAACATCATCAAAATTTGCGAAAAA

The match between the T1 sequence and the C1/C2 sequence is

TTTTACTCCTAATTACCAATTCTAACCATCCATATCACTTTTTGACGGACTC
CGTGAAAATAATTTTTGGCCAAATTTTCGCATTTTTTGTAAGGGGTAACAT
CAT

The match between the T2 sequence and the C1/C2 sequence is

[illegible]

.....

All of the human genome that has been fully sequenced by both the NIH-lead global sequencing project and the Celera Genomics, Inc. project. The gene descriptors for this chromosome do not yet exist. Without the positions and directions of the genes, it is not possible to select from among the possible connectrons to determine the real connectrons.

Human chromosome 22 has been processed and there 31,000 possible connectrons.


The gene descriptors for all the chromosomes of the human genome should become available within the year.

6. Permanent connectrons exist in prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes.

C1/C2 short loops are normally expressed as the 3'UTR of some gene. A class of connectron relationships exist that permit one C1/C2 short loop to control the existence of one or more T1-T2 long loops without being subject to any expression controls other than those of the gene to which the C1/C2 is 3'UTR. These connectron relationships are described as "permanent". Permanent connectrons exist in prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes.

Example of a prokaryote permanent connectron – E. coli

In this example the existence of the T1-T2 (3200-3210) long loop is controlled by a C1/C2 short loop (3432). The expression of this C1/C2 short loop is controlled only by the gene btuB.



A double stranded DNA loop of length 93.339 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3200. This T1 control element has the DNA sequence

AAGCGGCACTGCTCTTTAACAATTTATCAGACAATCTGTGTGGGCACTCG
AAGATACGGATTCTTAACGTCGCAAGACGAAAAATGAATACCAAGTCTCA
AGAGTGAACACGTAATTCATTACGAAGTTTAATTCTTTGAGCATCAAACCT
TTAAATTGAAGAGTTTGATCATGGCTCAGATTGAACGCTGGCGGCAGGCC
TAACACATGCAAGTCGAACGGTAACAGGAAACAGCTTGCTGTTTCGCTGA
CGAGTGGCGGACGGGTGAGTAATGTCTGGGAAACTGCCTGATGGAGGGG

GATAACTACTGGAAACGGTAGCTAATACCGCATAACGTCGCAAGACCAAA
GAGGGGGACCTTCGGGCCTCTTGCCATC

This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 3310. This T2 control element has the DNA sequence

CAGACAATCTGTGTGGGCACTCGAAGATACGGATTCTTAACGTCGCAAGA
CGAAAAATGAATACCAAGTCTCAAGAGTGAACACGTAATTCATTACGAAG
TTTAATTCTTTGAGCGTCAAACTTTTAAATTGAAGAGTTTGATCATGGCTC
AGATTGAACGCTGGCGGCAGGCCTAACACATGCAAGTCGAACGGTAACA
GGAAGAAGCTTGCTTCTTTGCTGACGAGTGGCGGACGGGTGAGTAATGTC
TGGGAAACTGCCTGATGGAGGGGGATAACTACTGGAAACGGTAGCTAAT
ACCGCATAACGTCGCAAGACCAAAGAGGGGGACCTTCGGGCCTCTTGCCA
TCGGATGTGCCCAGATGGGATTAGCTAGT

This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

rrsC	gltU	rrlC	rrfC	aspT	trpT	yifA	yifE	yifB
ilvL	ilvG_1	ilvM	ilvE	ilvD	ilvA	ilvY	ilvC	ppiC
b3776	rep	gppA	rhlB	trxA	rhoL	rho	rfe	wzzE
wecB	rffH	wecD	wecE	wzxE	yifM_2	wecG	yifK	
argX	hisR	leuT	proM	aslB	aslA	hemY	hemX	
hemD	cyaA	cyaY	b3808	dapF	uvrD	b3814	corA	
yigF	yigG	rarD	yigI	pldA	recQ	yigJ	yigK	pldb
yigL	yigM	metR	metE	ysgA	udp	yigN	ubiE	yigP
b3836	yigU	yigW_1	rfaH	yigC	ubiB	fadA	fadB	
pepQ	trkH	hemG						

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2
short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 3432 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene *btuB* and has the DNA sequence

5 TGC GCGGTCAGAAAATTATTTTAAATTCCTCTTGTCAGGCCGGAATAACT
 CCCTATAATGCGCCACCACTGACACGGAACAACGGCAAACACGCCGCCGG
 GTCAGCGGGGTTCTCCTGAGAACTCCGGCAGAGAAAGCAAAAATAAATG
 CTTGACTCTGTAGCGGGAAGGCGTATTATGCACACC...TGCAACTCGACTC
 CATGAAGTCGGAATCGCTAGTAATCGTGGATCAGAATGCCACGGTGAATA
 10 CGTTCCCGGGCCTTGTACACACCGCCCGTCACACCATGGGAGTGGGTTC
 AAAAGAAGTAGGTAGCTTAACCTTCGGGAGGGCGCTTACCACTTTGTGAT
 TCATGACTGGGGTGAAGTCGTAACAAGGTAACCGTAGGGGAACCTGCGGT
 TGGATCACCTCCTTACCTTAAAGAAGCGT

15 The match between the T1 sequence and the C1/C2 sequence is


AAGCGGCACTGCTCTTTAACAATTTATCAGACAATCTGTGTGGGCACTCG
 AAGATACGGATTCTTAACGTCGCAAGACGAAAAATGAATACCAAGTCTCA
 AGAGTGAACACGTAATTCATTACGAAGTTAATTCTTTGAGC

20 The match between the T2 sequence and the C1/C2 sequence is

CAGACAATCTGTGTGGGCACTCGAAGATACGGATTCTTAACGTCGCAAGA
 CGAAAAATGAATACCAAGTCTCAAGAGTGAACACGTAATTCATTACGAAG
 25 TTTAATTCTTTGAGCGTCAAACCTTTTAAATTGAAGAGTTTGATCATGGCTC
 AGATTGAACGCTGGCGGCAGGCCTAACACATGCAAGTCGAACGGTAACA
 GGAAGAAGCTTGCTTCTTTGCTGACGAGTGGCGGACGGGTGAGTAATGTC
 TGGGAAACTGCCTGATGGAGGGGGATAACTACTGGAAACGGTAGCTAAT
 ACCGCATAACGTCGCAAGACCAAAGAGGGGGACCTTCGGGCCTCTTGCCA
 30 TCGGATGTGCCCAGATGGGATTAGCTAGT

Example of an archea permanent connectron – *H. pylori*

In this example the existence of the T1-T2 (812-882) long loop is controlled by a C1/C2 short loop (1241). The expression of this C1/C2 short loop is controlled only by the gene HP1535.



A double stranded DNA loop of length 96.385 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 812. This T1 control element has the DNA sequence

TTTACTCATAGGGTTTTATAGTTCCTAGCGGAACTAAAGCA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 882. This T2 control element has the DNA sequence

TAGCGGAACTAAAGCATTCATCCCAAACACTAAAGATATTTGG

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

HP0999	HP1000	HP1001	HP1002	HP1003	HP1005	HP1006
HP1008	HP1009	HPtRNA-Pro	HP1010	HP1011	HP1013	HP1015
HP1017	HP1018	HP1020	HP1021	HP1022	HP1023	HP1024
HP1025	HP1027	HP1028	HP1030	HP1031	HP1033	HP1034
HP1038	HP1039	HP1040	HP1041	HP1042	HP1043	HP1044

HP1045 HP1046 HP1051 HP1052 HP1055 HP1056 HP1058
 HP1060 HP1065 HPtRNA-Ser HP1066 HP1067 HP1069 HP1070
 HP1074 HP1075 HP1076 HP1077 HP1078 HP1079 HP1080
 HP1081 HP1083 HP1084 HP1085 HP1088 HP1091 HP1092
 5 HP1093 HP1094 HP1095 HP1096

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

10 A C1/C2 short loop on chromosome 1 whose identifier is 1241 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene HP1535 and has the DNA sequence

15 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCATTTCATCCC
 AAACA

The match between the T1 sequence and the C1/C2 sequence is


20 TTTTACTCATAGGGTTTTTATAGTTCCTAGCGGAACTAAAGCA

The match between the T2 sequence and the C1/C2 sequence is

25 TAGCGGAACTAAAGCATTTCATCCCAAACA

Example of a single-celled permanent connectron – *S. cerevisiae*

30 In this example the existence of the T1-T2 (5515-5533) long loop is controlled by a C1/C2 short loop (6102). The expression of this C1/C2 short loop is controlled only by the gene YNL339C.



A double stranded DNA loop of length 6.466 kilo-bases on chromosome 12 is bounded on the left by a T1 sequence whose identifier is 5515. This T1 control element has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTGATTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 5533. This T2 control element has the DNA sequence

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCAGTGTGTTGATTAGTGTGTTGTTGC
ACGGCAGTAGCGAGAGACAAGTGGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

YLR467W

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 14 whose identifier is 6102 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YNL339C and has the DNA sequence

AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTGAACATCCGGGTAAGAGACAACAGGGCT

The match between the T1 sequence and the C1/C2 sequence is


AGGAAATTGTTGTTACGAAAGTCAGTGATTATGTATTGTGTAGTATAGTAT
ATTGTAAGAAATTTTTTTTTTCTAGGGAATATGCGTTTTGATGTAGTAGTAT
TTCAGTGTGTTTATTAGTGTGTTGTTGCACGGCAGTAGCGAGAGACAAGTG
GGAAAGAGTAGGATAAAAAGACAATCTATAAAAAGTAAACATAAAATAA
AGGTAGTAAGTAGCTTTTGGTTG

The match between the T2 sequence and the C1/C2 sequence is

ATTATGTATTGTGTAGTATAGTATATTGTAAGAAATTTTTTTTTTCTAGGGA
ATATGCGTTTTGATGTAGTAGTATTTCACTGTTTGTGTTGTTGTTGC
ACGGCAGTAGCGAGAGACAAGTGGAAAGAGTAGGATAAAAAGACAATC
TATAAAAAGTAAACATAAAATAAAGGTAGTAAGTAGCTTTTGGTTGAACA
TCCGGGTAAGAGACAACAGGGCT

Example of a multi-celled permanent connectron – *C. elegans*

In this example the existence of the T1-T2 (5515-5533) long loop is controlled by a C1/C2 short loop (6102). The expression of this C1/C2 short loop is controlled only by the gene YNL339C.



A double stranded DNA loop of length 30.606 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 569. This T1 control element has the DNA sequence

AAATCGAGCCCGTAAATCGACACAAGCGCTACAGTAGTC

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 596. This T2 control element has the DNA sequence

AGTGCTACAGTAGTCATTTAAAGAATTACTGTAGTTTTTCGCT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 5 whose identifier is 24442 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene F20D6.4 and has the DNA sequence

GAGCCCGTAAATCGACACAAGCGCTACAGTAGTCATTTAAAGAATTACTG
TAGTTTTTC


The match between the T1 sequence and the C1/C2 sequence is

7. Transient connectrons exist in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

A class of connectron relationships exist that permit one C1/C2 short loop to control the existence of one or more T1-T2 long loops such that this C1/C2 short loop is itself subject to expression control by another T1-T2 long loop which surrounds it. These connectron relationships are described as “transient”. Transient connectrons exist in prokaryotes, archaea, single-celled eukaryotes and multi-celled eukaryotes.

Example of a prokaryote transient connectron – E. coli

In this example the existence of the T1-T2 (3227-3329) long loop is controlled by the C1/C2 (3225) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (3216-3224) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (3223) short loop. The C1/C2 (3225) short loop is the transient connectron.



A double stranded DNA loop of length 93.464 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 3216. This T1 control element has the DNA sequence

AGCGCAAGCGAAGCTCTTGATCGAAGCCCCGGTAAACGGCGGCCGTAAC
ATAACGGTCCTAAGGTAGCGAAATTCCTTGTCGGGTAAGTTCCGACCTGC
ACGAATGGCGTAATGATGGCCAGGCTGTCTCCACCCGAGACTCAGTGA
TTGAACTCGCTGTGAAGATGCAGTGTACCCGCGGCAAGACGGAAAGACCC
5 CGTGAACCTTTACTATAGCTTGACACTGAACATTGAGCCTTGATGTGTAGG
ATAGGTGGGAGGCTTTGAAGTGTGGACGCCAGTCTGCATGGAGCCGACCT
TGAAATACCACCCTTTAATGTTTGATGTTCTAACGT

This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 3324. This T2 control element has the DNA sequence

CCCGGTAAACGGCGGCCGTAACATAACGGTCCTAAGGTAGCGAAATTC
TTGTCGGGTAAGTTCCGACCTGCACGAATGGCGTAATGATGGCCAGGCTG
TCTCCACCCGAGACTCAGTGAATTTGAACTCGCTGTGAAGATGCAGTGTA
15 CCCGCGGCAAGACGGAAAGACCCCGTGAACCTTTACTATAGCTTGACACT
GAACATTGAGCCTTGATGTGTAGGATAGGTGGGAGGCTTTGAAGTGTGGA
CGCCAGTCTGCATGGAGCCGACCTTGAAATACCACCCTTTAATGTTTGATG
TTCTAACGTTGACCCGTAATCCGGGTTGCGGACAGT

This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

rrfC	aspT	trpT	yifA	yifE	yifB	ilvL	ilvG_1	ilvM
ilvE	ilvD	ilvA	ilvY	ilvC	ppiC	b3776	rep	gppA
25 rhlB	trxA	rhoL	rho	rfe	wzzE	wecB	rffH	wecD
wecE	wzxE	yifM_2	wecG	yifK	argX	hisR	leuT	
proM	aslB	aslA	hemY	hemX	hemD	cyaA	cyaY	
b3808	dapF	uvrD	b3814	corA	yigF	yigG	rarD	yigI
pldA	recQ	yigJ	yigK	pldB	yigL	yigM	metR	metE
30 ysgA	udp	yigN	ubiE	yigP	b3836	yigU	yigW_1	rfaH
yigC	ubiB	fadA	fadB	pepQ	trkH	hemG	rrsA	ileT
rrlA								

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

5 A C1/C2 short loop on chromosome 1 whose identifier is 3225 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrlC and has the DNA sequence

10 AAACAGAATTTGCCTGGCGGCCGTAGCGCGGTGGTCCCACCTGACCCCAT
GCCGAAGTCTCAGAAAGTCAAACGCGGTAGCGCCGATGGTAGTGTGGGGTCTC
CCCATGCGAGAGTAGGGAAGTCCAGGCATCAAATTA

15 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 3323 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrlA and has the DNA sequence

20 GCGAAGCTCTTGATCGAAGCCCCGGTAAACGGCGGCCGTAAGTATAACGG
TCCTAAGGTAGCGAAATTCCTTGTCTGGGTAAAGTTCGACCTGCACGAATG
GCGTAATGATGGCCAGGCTGTCTCCACCCGAGACTCAGTGAAATTGAACT
CGCTGTGAAGATGCAGTGTACCCGCGGCAAGACGGA...AACAGAATTTGC
25 CTGGCGGCAGTAGCGCGGTGGTCCCACCTGACCCCATGCCGAAGTCTCAGAA
GTGAAACGCGGTAGCGCCGATGGTAGTGTGGGGTCTC

The match between the T1 sequence and the C1/C2 sequence is

30 GCGAAGCTCTTGATCGAAGCCCCGGTAAACGGCGGCCGTAAGTATAACGG
TCCTAAGGTAGCGAAATTCCTTGTCTGGGTAAAGTTCGACCTGCACGAATG
GCGTAATGATGGCCAGGCTGTCTCCACCCGAGACTCAGTGAAATTGAACT

CGCTGTGAAGATGCAGTGTACCCGCGGCAAGACGGAAAGACCCCGTGAA
 CCTTTACTATAGCTTGACACTGAACATTGAGCCTTGATGTGTAGGATAGGT
 GGGAGGCTTTGAAGTGTGGACGCCAGTCTGCATGGAGCCGACCTTGAAAT
 ACCACCCTTTAATGTTTGATGTTCTAACGT

5

The match between the T2 sequence and the C1/C2 sequence is

CCCGGTAAACGGCGGCCGTAACATAACGGTCCTAAGGTAGCGAAATTCC
 TTGTCGGGTAAGTTCCGACCTGCACGAATGGCGTAATGATGGCCAGGCTG
 TCTCCACCCGAGACTCAGTGAAATTGAACTCGCTGTGAAGATGCAGTGTA
 CCCGCGGCAAGACGGAAAGACCCCGTGAACCTTTACTATAGCTTGACACT
 GAACATTGAGCCTTGATGTGTAGGATAGGTGGGAGGCTTTGAAGTGTGGA
 CGCCAGTCTGCATGGAGCCGACCTTGAAATACCACCCTTTAATGTTTGATG
 TTCTAACGTTGACCCGTAATCCGGGTTGCGGACAGT

10

15

A double stranded DNA loop of length 93.749 kilo-bases on chromosome 1 is
 bounded on the left by a T1 sequence whose identifier is 3227. This T1 control
 element has the DNA sequence

20

AGCGCCGATGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGGAACTGCCA
 GG

25

This double stranded DNA loop is bounded on the right by a T2 control element
 whose identifier is 3329. This T2 control element has the DNA sequence

CATGCGAGAGTAGGGAACTGCCAGGCATCAAATAAAACGAAAGGCTCAG
 TCG

30

This long T1/T2 double stranded DNA loop modulates the expression of the
 following genes

aspT trpT yifA yifE yifB ilvL ilvG_1 ilvM ilvE
 ilvD ilvA ilvY ilvC ppiC b3776 rep gppA rhlB
 trxA rhoL rho rfe wzzE wecB rffH wecD wecE
 5 wzxE yifM_2 wecG yifK argX hisR leuT proM
 aslB aslA hemY hemX hemD cyaA cyaY b3808
 dapF uvrD b3814 corA yigF yigG rarD yigI pldA
 recQ yigJ yigK pldB yigL yigM metR metE ysgA
 udp yigN ubiE yigP b3836 yigU yigW_1 rfaH yigC
 10 ubiB fadA fadB pepQ trkH hemG rrsA ileT rrlA
 rrfA

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 3225 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene rrlC and has the DNA sequence

AAACAGAATTTGCCTGGCGGCCGTAGCGCGGTGGTCCCACCTGACCCCAT
 GCCGAAGTCTCAGAAAGTCAAACGCGGTAGCGCCGATGGTAGTGTGGGGTCTC
 CCCATGCGAGAGTAGGGAACTGCCAGGCATCAAATTA

The match between the T1 sequence and the C1/C2 sequence is

AGCGCCGATGGTAGTGTGGGGTCTCCCCATGCGAGAGTAGGGAACTGCCA
 GG


The match between the T2 sequence and the C1/C2 sequence is

CATGCGAGAGTAGGGAACTGCCAGGCATCAAAT


Example of an archea transient connectron – *M. jannaschii*

5 In this example the existence of the T1-T2 (1139-1159) long loop is controlled by the C1/C2 (533) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (532-622) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (1629) short loop. The C1/C2 (533) short loop is the transient connectron.

10



15



20

25

A double stranded DNA loop of length 78.672 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 532. This T1 control element has the DNA sequence

30

ATATGTTTGAAATTTGAAAATAAGAGTATTTAG

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 622. This T2 control element has the DNA sequence

35

TTGAAAATAAGAGCATTAGAAAGTTATTAATTAGTTCAAAGGATTTT

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

	MJ0486	MJ0487	MJ0488	MJ0489	MJ0490	MJ0492	MJ0493
5	MJ0494	MJ0495	MJ0496	MJ0497	MJ0499	MJ0500	MJ0501
	MJ0502	MJ0503	MJ0504	MJ0506	MJ0507	MJ0508	MJ0509
	MJ0510	MJ0511	MJ0512	MJ0513	MJ0514	MJ0514	MJ0517
	MJ0519	MJ0520	MJ0521	MJ0522	MJ0523	MJ0525	MJ0526
	MJ0526	MJ0529	MJ0530	MJ0531	MJ0532	MJ0534	MJ0535
10	MJ0536	MJ0538	MJ0539	MJ0540	MJ0541	MJ0542	MJ0543
	MJ0544	MJ0545	MJ0547	MJ0548	MJ0549	MJ0550	MJ0552
	MJ0553	MJ0554	MJ0555	MJ0556	MJ0557	MJ0558	MJ0559
	MJ0560	MJ0561	MJ0562	MJ0563	MJ0564	MJ0565	MJ0566
	MJ0568	MJ0569	MJ0570				

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 1 whose identifier is 533 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ0485 and has the DNA sequence

ATTTTATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGA
GTTTATTGAATT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 1629 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1597 and has the DNA sequence

ATATGTTTGAAATTTGAAAATAAGAGTATTTAGAAGTTATTAATTAGTTCA
AAGGATTTTTATTTAATTTCTAAGGGTTTGCTGGTTTGATTATTTAGAATAT
TTGAGTTTATTGAATTATTCAGATTTTAAAAATTA

5

The match between the T1 sequence and the C1/C2 sequence is

ATATGTTTGAAATTTGAAAATAAGAGTATTTAG

10

The match between the T2 sequence and the C1/C2 sequence is

ATTTAGAAGTTATTAATTAGTTCAAAGGATTTT

15

A double stranded DNA loop of length 14.509 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1139. This T1 control element has the DNA sequence

20

ATTTATTAATTAGTTCAAAGGATTTTATTTAATTTCTAAGGGTTAGCTGG
TTTGATTGTTTAAAATATTTGAGTTTA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1159. This T2 control element has the DNA sequence

25

ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTTAGAATATTTGAGTTTAT
TGAATTATTCAGATTTTAAAAATTA

30

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

MJ1096 MJ1097 tRNA-Arg-3 MJ1098 MJ1099 MJ1100 MJ1101
 MJ1102 MJ1103 MJ1104 MJ1105 MJ1106 MJ1107 MJ1108

5 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 533 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ0485 and has the DNA sequence

10 ATTTTATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTAGAAATATTTGAGTTTATTGAATT

The match between the T1 sequence and the C1/C2 sequence is


15 ATTTTATTTAATTTCTAAGGGTTAGCTGGTTTGATT

The match between the T2 sequence and the C1/C2 sequence is

20 ATTTAATTTCTAAGGGTTAGCTGGTTTGATTATTAGAAATATTTGAGTTTATTGAATT

25 Example of a single-celled transient connectron – *S. cerevisiae*

In this example the existence of the T1-T2 (2840-2859) long loop is controlled by the C1/C2 (298) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (293-320) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (86) short loop. The C1/C2 (298) short loop is the transient connectron.



A double stranded DNA loop of length 38.470 kilo-bases on chromosome 2 is bounded on the left by a T1 sequence whose identifier is 293. This T1 control element has the DNA sequence

GAATTGTTGGAATAAAAATCCACTATCGTCTATCAACTAATAGTTATATTA
TCAATATATTATCATATACGGTGTTAAGATGATGACATAAGTTATGAGAA
GCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCAAGGATTGATAATGTAAT
AGGATAATGAAACATATAAAACGGAATGAGGAATAATCGTAATATTAGT
ATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTTGAGGAGAAC
TTCTAGT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 320. This T2 control element has the DNA sequence

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTC
GAGGAGAACTTCTAGTATATTCTGTA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

YBL005W-B TS(AGA)B YBL004W YBL003C YBL002W YBL001C
YBR001C YBR002C YBR003W YBR004C YBR005W YBR006W
YBR007C YBR008C YBR009C YBR010W YBR011C YBR012C

5

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

10

A C1/C2 short loop on chromosome 2 whose identifier is 298 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YBL005W-B and has the DNA sequence

15

ATCTATTACATTATGGGTGGTATGTTGGAATAAAAATCCACTATCGTCTAT
CAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGATGAT
GACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCA
AGGATTGATAATGTAATAGGATAATGAAACATATAAAACGGAATGAGGA
ATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCCT
ATATCCTTGAGGAGAACTTCTAGTATATTCTGTATACCTAATATTATAGCC
TTTATCAACAATGGAATCCCAACAATTATCTCAACATTC

20

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25

A C1/C2 short loop on chromosome 1 whose identifier is 86 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YAR009C and has the DNA sequence

30

ATCTATTACATTATGGGTGGTATGTTGGAATAGAAATCAACTATCATCTAC
TAACTAGTATTTACATTACTAGTATATTATCATATACGGTGTTAGAAGATG
ACGCAAATGATGAGAAATAGTCATCTAAATTAGTGGAAGCTGAAACGCA
AGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACGGAAT

GAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGG
ATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAATATT
ATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCACCCAT
TTCTCAGAA

5

The match between the T1 sequence and the C1/C2 sequence is

AAACATATAAAACGGAATGAGGAATAATCGTAATATTAGTATGTAGAAAT
ATAGATTCCATTTTGAGGATTCCTATATCCT

10

The match between the T2 sequence and the C1/C2 sequence is

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTC
GAGGAGAACTTCTAGTATATTCTGTA

15

A double stranded DNA loop of length 5.302 kilo-bases on chromosome 7 is bounded
on the left by a T1 sequence whose identifier is 2840. This T1 control element has
the DNA sequence

20

TCTGTTGGAATAAAAATCCACTATCGTCTATCAACTAATAGTTATATTATC
AATATATTATCATATACGGTGTTAAGATGATGACATAAGTTATGAGAAGC
TGTCATCGAAGTTAGAGGAAGCTGAAACGCAAGGATTGATAATGTAATAG
GATCAATGAATATAAACATATAAAACGGAATGAGGAATAATCGTAATATT
AGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTCGAGGAG
AACTTCTAGTATATTCTGTATACCTAAATTATAGCCTTTATCAACAATGGA
ATCCCAACAA

25

30

This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 2859. This T2 control element has the DNA sequence

CTATCAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGA
 TGATGACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAA
 CGCAAGGATTGATAATGTAATAGGATCAATGAATATAAACATATAAAACG
 GAATGAGGAATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTT
 5 GAGGATTCCTATATCCTCGAGGAGAACTTCTAGTATATTCTGTATACCTAA
 TATTATAGCCTTTATCAACAATGGAATCCCAACAATTATCTCAACATTCAC
 ATATTTCTCAT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2
 10 short loops.

A C1/C2 short loop on chromosome 2 whose identifier is 298 controls the expression
 of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA
 single strand that is 3'UTR to the gene YBL005W-B and has the DNA sequence

ATCTATTACATTATGGGTGGTATGTTGGAATAAAAAATCCACTATCGTCTAT
 CAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGATGAT
 GACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCA
 AGGATTGATAATGTAATAGGATAATGAAACATATAAAACGGAATGAGGA
 15 ATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCT
 ATATCCTTGAGGAGAACTTCTAGTATATTCTGTATACCTAATATTATAGCC
 TTTATCAACAATGGAATCCCAACAATTATCTCAACATTC
 20

The match between the T1 sequence and the C1/C2 sequence is

TGTTGGAATAAAAAATCCACTATCGTCTATCAACTAATAGTTATATTATCAA
 TATATTATCATATACGGTGTTAAGATGATGACATAAGTTATGAGAAGCTG
 TCATCGAAGTTAGAGGAAGCTGAA
 25

The match between the T2 sequence and the C1/C2 sequence is

30

CTATCAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGA
TGATGACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAA

5

Example of a multi-celled transient connectron – *C. elegans*


In this example the existence of the T1-T2 (22072-22108) long loop is controlled by the C1/C2 (125) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (110-129) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (16859) short loop. The C1/C2 (125) short loop is the transient connectron.

10

15

20

25



30

A double stranded DNA loop of length 18.855 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 110. This T1 control element has the DNA sequence

AGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGC

35

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 129. This T2 control element has the DNA sequence

TTCTCCCGCATTTTTTGTAGATCTACGTAGATCAAACCGAAATGAGGCACT
TTCTGAATCCACGAGCTAGGCTTAAGCTTAGGCTTAAGCTTAGGCCTTTTC
TCAGGCTTAGGCTTAGGCTTA

5

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

ZC123.3 ZC123.2

10

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 1 whose identifier is 125 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.3 and has the DNA sequence

15

ACGCGCCGTAAATCTACCCCAGATATGGCCGAGCCAAAATGGCCTAGTTC
GGCAAACCTCTTTCATTTCAATTTATGAGGGAAGCCAGAA

20

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25

A C1/C2 short loop on chromosome 4 whose identifier is 16859 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene F58E2.7 and has the DNA sequence

CTTAGGCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTT
AGGCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTTAG
GCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGCT
TAAGCTTAGACTTA

30

The match between the T1 sequence and the C1/C2 sequence is

AGCTTAGGCTTAAGCTTAGGCTTAAGCTTAGGC

The match between the T2 sequence and the C1/C2 sequence is

TAGGCTTAAGCTTAGGCTTAAGCTTAGGC

A double stranded DNA loop of length 51.031 kilo-bases on chromosome 5 is bounded on the left by a T1 sequence whose identifier is 22072. This T1 control element has the DNA sequence

CGCAACGCGCCGTAAATCTACCCCAGATATGGCCGAGCCAAAATGACCTA
GTTCGGC

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 22108. This T2 control element has the DNA sequence

TGACAATCGCCTGCCGGACAACGCGTGGAAAAGTGTCGTGTACTCCACAC
GGACAAATACATTTAGTTTTACAATAAAATCGAACCGCGACGCGACACG
CAACGCGACGTAAATCTACCCCAGATATGGCCGAGCCAAAATGGCCTAGT
TCGGCAAACCTTCTATTTC

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

F36H9.3 F36H9.4 F36H9.5 F36H9.2 F36H9.1 F36H9.6

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 125 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZC123.3 and has the DNA sequence

ACGCGCCGTAAATCTACCCCAGATATGGCCGAGCCAAAATGGCCTAGTTC
GGCAAACCTCTTTCATTTCAATTTATGAGGGAAGCCAGAA

The match between the T1 sequence and the C1/C2 sequence is

ACGCGCCGTAAATCTACCCCAGATATGGCCGAGCCAAAATG

The match between the T2 sequence and the C1/C2 sequence is


CGTAAATCTACCCCAGATATGGCCGAGCCAAAATGGCCTAGTTCGGCAAA
CTCTT

8. Self-limiting connectrons occur in prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes

A class of connectron relationships exist that permit one C1/C2 short loop to control the existence of the T1-T2 long loop that surrounds it. These connectron relationships are described as “self-limiting”. Self-limiting connectrons exist in prokaryotes, archea, single-celled eukaryotes and multi-celled eukaryotes.

Example of a prokaryotic self-limiting connectrons – E. coli

In this example the existence of the T1-T2 (1704-1718) long loop is controlled by two C1/C2 (1705 and 1713) short loops. The expression of these C1/C2 short loops is controlled by the existence of the T1-T2 (1704-1718) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the two C1/C2 (1705 and 1713) short loops. The C1/C2 (1705 and 1713) short loops are the self-limiting connectrons.



A double stranded DNA loop of length 15.259 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1704. This T1 control element has the DNA sequence

CGCCCCGTTACACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACT
GTTAATCCGTATGTCACTGGT

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1718. This T2 control element has the DNA sequence

5 TTCAGTCGGTAGAACGGCGGACTGTTAATCCGTATGTCACTGGTTCGAGTC
CAGTCAGAGGAGCCAAATTC

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

10 asnT b1978 b1979 b1980 shiA amn b1983 asnW
yeeO asnU

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15 A C1/C2 short loop on chromosome 1 whose identifier is 1705 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene and has the DNA sequence

20 CGCCCCGTTACACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACT
GTTAATCCGTATGTCACTGGTTCGAGTCCAGTCAGAGGAGCCAAATTC

25 A C1/C2 short loop on chromosome 1 whose identifier is 1713 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene asnW and has the DNA sequence

30 CACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACTGTTAATCCGTAT
GTCACTGGTTCGAGTCCAGTCAGAGGAGCCAAATT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 1705 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene and has the DNA sequence

CGCCCCGTTACACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACT
GTTAATCCGTATGTCACTGGTTCGAGTCCAGTCAGAGGAGCCAAATTC

The match between the T1 sequence and the C1/C2 sequence is

CGCCCCGTTACACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACT
GTTAATCCGTATGTCACTGGT

The match between the T2 sequence and the C1/C2 sequence is

TTCAGTCGGTAGAACGGCGGACTGTTAATCCGTATGTCACTGGTTCGAGTC
CAGTCAGAGGAGCCAAATTC

A C1/C2 short loop on chromosome 1 whose identifier is 1713 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene asnW and has the DNA sequence

CACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACTGTTAATCCGTAT
GTCACTGGTTCGAGTCCAGTCAGAGGAGCCAAATT

The match between the T1 sequence and the C1/C2 sequence is

CACGATTCCTCTGTAGTTCAGTCGGTAGAACGGCGGACTGTTAATCCGTAT
GTCACTGGT


The match between the T2 sequence and the C1/C2 sequence is

TTCAGTCGGTAGAACGGCGGACTGTTAATCCGTATGTCACTGGTTCGAGTC
CAGTCAGAGGAGCCAAATT

5

Example of a archea self-limiting connectrons – *M. jannaschii*

In this example the existence of the T1-T2 (1447-1471) long loop is controlled by two C1/C2 (1448 and 1470) short loops. The expression of these C1/C2 short loops is controlled by the existence of the T1-T2 (1447-1471) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the two C1/C2 (1705 and 1713) short loops. The C1/C2 (1448 and 1470) short loops are the self-limiting connectrons.



A double stranded DNA loop of length 22.675 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 1447. This T1 control element has the DNA sequence

TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTC
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTTATGATA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1471. This T2 control element has the DNA sequence

CAACTAACAACCGTATCGAATTTACCATTACTTGGAATCTATTTAAAACC
TCTTTAATCTTGTGATAATAAATTCTAATCGATTTCGTGACTTAT

5 This long T1/T2 double stranded DNA loop modulates the expression of the following genes

MJ1402 MJ1403 MJ1404 MJ1405 MJ1406 MJ1407 MJ1408
MJ1409 MJ1410 MJ1411 MJ1412 MJ1413 MJ1414 MJ1415
10 MJ1416 MJ1417 MJ1418 MJ1419 MJ1420

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15 A C1/C2 short loop on chromosome 1 whose identifier is 1448 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1401 and has the DNA sequence

20 TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTTATGATAATAAATT
CTAATCGATTTCGTGACTTAT

25 A C1/C2 short loop on chromosome 1 whose identifier is 1470 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1420 and has the DNA sequence

30 TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTTGTGATAATAAATT
CTAATCGATTTCGTG

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 1470 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1420 and has the DNA sequence

TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTTGTGATAATAAATT
CTAATCGATTTCGTG

The match between the T1 sequence and the C1/C2 sequence is

TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTT

The match between the T2 sequence and the C1/C2 sequence is

CAACTAACAACCGTATCGAATTTACCATTACTTGGAATCTATTTAAAACC
TCTTTAATCTTGTGATAATAAATTCTAATCGATTTCGTG

A C1/C2 short loop on chromosome 1 whose identifier is 1448 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene MJ1401 and has the DNA sequence

TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAATCTATTTAAAACCTCTTTAATCTTATGATAATAAATT
CTAATCGATTTCGTGACTTAT

The match between the T1 sequence and the C1/C2 sequence is


TTATAGAACATTATGAAGCTTTTTACTCAACTAACAACCGTATCGAATTTA
CCATTACTTGGAAATCTATTTAAAACCTCTTTAATCTTATGATA

The match between the T2 sequence and the C1/C2 sequence is

CAACTAACAACCGTATCGAATTTACCATTACTTGGAAATCTATTTAAAACC
TCTTTAATCTT

Example of a single-celled self-limiting connectron – *S. cerevisiae*

In this example the existence of the T1-T2 (293-320) long loop is controlled by C1/C2 (298) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (293-320) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (298) short loop. The C1/C2 (298) short loop is the self-limiting connectron.



A double stranded DNA loop of length 38.470 kilo-bases on chromosome 2 is bounded on the left by a T1 sequence whose identifier is 293. This T1 control element has the DNA sequence

GAATTGTTGGAATAAAAATCCACTATCGTCTATCAACTAATAGTTATATTA
TCAATATATTATCATATACGGTGTTAAGATGATGACATAAGTTATGAGAA
GCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCAAGGATTGATAATGTAAT

AGGATAATGAAACATATAAAACGGAATGAGGAATAATCGTAATATTAGT
ATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTTGAGGAGAAC
TTCTAGT

5 This double stranded DNA loop is bounded on the right by a T2 control element
whose identifier is 320. This T2 control element has the DNA sequence

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCTC
GAGGAGAACTTCTAGTATATTCTGTA

10

This long T1/T2 double stranded DNA loop modulates the expression of the
following genes

YBL005W-B TS(AGA)B YBL004W YBL003C YBL002W YBL001C
15 YBR001C YBR002C YBR003W YBR004C YBR005W YBR006W
YBR007C YBR008C YBR009C YBR010W YBR011C YBR012C

This long T1/T2 double stranded DNA loop modulates the expression of the
following C1/C2 short loops

20

A C1/C2 short loop on chromosome 2 whose identifier is 298 controls the expression
of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is
expressed as a RNA single strand that is 3'UTR to the gene YBL005W-B and has the
DNA sequence

25

ATCTATTACATTATGGGTGGTATGTTGGAATAAAAATCCACTATCGTCTAT
CAACTAATAGTTATATTATCAATATATTATCATATACGGTGTTAAGATGAT
GACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCA
AGGATTGATAATGTAATAGGATAATGAAACATATAAAACGGAATGAGGA
30 ATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCT
ATATCCTTGAGGAGAACTTCTAGTATATTCTGTATACCTAATATTATAGCC
TTTATCAACAATGGAATCCCAACAATTATCTCAACATTC

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

- 5 A C1/C2 short loop on chromosome 2 whose identifier is 298 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YBL005W-B and has the DNA sequence

10 ATCTATTACATTATGGGTGGTATGTTGGAATAAAAATCCACTATCGTCTAT
CAACTAATAGTTATATTATCAATATATTATCATATACGGTGTAAAGATGAT
GACATAAGTTATGAGAAGCTGTCATCGAAGTTAGAGGAAGCTGAAGTGCA
AGGATTGATAATGTAATAGGATAATGAAACATATAAAACGGAATGAGGA
ATAATCGTAATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCT
15 ATATCCTTGAGGAGAACTTCTAGTATATTCTGTATACCTAATATTATAGCC
TTTATCAACAATGGAATCCCAACAATTATCTCAACATTC

The match between the T1 sequence and the C1/C2 sequence is

20 TGTTGGAATAAAAATCCACTATCGTCTATCAACTAATAGTTATATTATCAA
TATATTATCATATACGGTGTAAAGATGATGACATAAGTTATGAGAAGCTG
TCATCGAAGTTAGAGGAAGCTGAAGTGCAAGGATTGATAATGTAATAGGA
TAATGAAACATATAAAACGGAATGAGGAATAATCGTAATATTAGTATGTA
GAAATATAGATTCCATTTTGAGGATTCCTATATCCTTGAGGAGAACTTCTA
GT

25


The match between the T2 sequence and the C1/C2 sequence is

AATATTAGTATGTAGAAATATAGATTCCATTTTGAGGATTCCTATATCCT

30

Example of a multi-celled self-limiting connectron – C. elegans

In this example the existence of the T1-T2 (293-320) long loop is controlled by C1/C2 (298) short loop. The expression of this C1/C2 short loop is controlled by the existence of the T1-T2 (293-320) long loop. The existence of this T1-T2 long loop is itself determined by the expression of the C1/C2 (298) short loop. The C1/C2 (298) short loop is the self-limiting connectron.



A double stranded DNA loop of length 89.919 kilo-bases on chromosome 4 is bounded on the left by a T1 sequence whose identifier is 17154. This T1 control element has the DNA sequence

AAATTTCCGGCAAATCGGCAAACACTGGCAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 17190. This T2 control element has the DNA sequence

AATTTGCCGATTTGCCGAATTTGTCGACA

This long T1/T2 double stranded DNA loop modulates the expression of the following genes

R08C7.11 M01H9.2 M01H9.3 M01H9.4 M01H9.1 ZK180.1 ZK180.2
ZK180.3 ZK180.4 ZK180.5 ZK180.6 ZK185.3 ZK185.2

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 4 whose identifier is 17155 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene R08C7.1 and has the DNA sequence

AAATTTCCGGCAAATCGGCAAACCTGGCAATTTGCCGATTTGCCGAATTTGT
CGACA

A C1/C2 short loop on chromosome 4 whose identifier is 17171 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene ZK180.2 and has the DNA sequence

TGGAAATTTTCAGAATTTCAATTTTAATCGGCAAATTTGTACGCATCCTATG
AATTT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 4 whose identifier is 17155 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene R08C7.1 and has the DNA sequence

AAATTTCCGGCAAATCGGCAAACCTGGCAATTTGCCGATTTGCCGAATTTGT
CGACA

The match between the T1 sequence and the C1/C2 sequence is

AAATTTCCGGCAAATCGGCAAACCTGGCAA

The match between the T2 sequence and the C1/C2 sequence is

AATTTGCCGATTTGCCGAATTTGTCGACA

5


09865925 053001

9. Geneless connectrons exist in single-celled and multi-celled eukaryotes

Normally T1-T2 long loops contain genes whose expression is regulated by the existence of the long loop. When a T1-T2 long loop does not contain any genes it is described as being “geneless”. The existence of the T1-T2 long loop is itself controlled by one or more C1/C2 short loops that may be on the same or different chromosomes. The geneless T1-T2 long loops must contain one or more C1/C2 short loops.

Example of a single-celled geneless connectron – *S. cerevisiae*

In this example the existence of the T1-T2 (1537-1559) long loop is controlled by three C1/C2 (3789, 5289 and 5753) short loops. The expression of 21 C1/C2 (1538 through 1558) short loops are controlled by the existence of the T1-T2 (1537-1559) long loop.



A double stranded DNA loop of length 4.825 kilo-bases on chromosome 4 is bounded on the left by a T1 sequence whose identifier is 1537. This T1 control element has the DNA sequence

ATGAGATATATGTGGGTAATTAGATAATTGTTGGGATTCCATTGTTGATAA
AGGCTATAATATTAGGTATACAGAATATACTAGAAGTTCTCCTCGAGGAT
TTAGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTATAAATAT
TATTATCATCGTTTTATATGTTAATATTCATTGATCCTATTACATTATCAAT

CCTTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGT
CATCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATACTAGTTA
GTAGATGATAGTTGATTTTTATTCCAACATACCACCCATAATGTAATAGAT
CTAAT

5

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1559. This T2 control element has the DNA sequence

ATGAGATATATGTGGGTAATTAGATAATTGTTGGGATTCCATTGTTGATAA
AGGCTATAATATTAGGTATACAGAATATACTAGAAGTTCTCCTCGAGGAT
TTAGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTATAAATAT
TATTATCATCGTTTTATATGTTAATATTCATTGATCCTATTACATTATCAAT
CCTTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGT
CATCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATACTAGTTA
GTAGATGATAGTTGATTTTTATTCCAACATACCACCCATAATGTAATAGAT
CTAAT

10

15

There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 4 whose identifier is 1538 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

ATGAGATATATGTGGGTAATTAGATAATTGTTGGGATTCCATTGTTGATAA
AGGCTATAATATTAGGTATACAGAATATACTAGAAGTTCTCCTCGAGGAT
TTAGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTATAAATAT
TATTATCATCGTTTTATATGTTAATATTCATTGATCCTATTACATTATCAAT
CCTTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGT
CATCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATACTAGTTA

30

GTAGATGATAGTTGATTTTTATTCCAACATACCACCCATAATGTAATAGAT
CTAATGAATCCATTTGTTTGTTAATAGTTT

This T1-T2 loop also modulates the C1/C2 short loops numbered 1539 to 1557

A C1/C2 short loop on chromosome 4 whose identifier is 1558 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

AGCTTCTCATAACTTATGTCATCATCTTAACACCGTATATGATAATATATT
GATAATATAACTTGTTGGAATAAAAATCAACTATCATCTACTAACTAGTAT
TTACGTTACTAGTATATTATCATATACGGTGTTAGAAGATGACGCAAATG
ATGAGAAATAGTCATCTAAATTAGTGGAAGCTGA...GTCTATCTGGCGAAT
ATAAATTTTTACGCTACACACGTCATCGACATCTAAATATGACAGTCGCTG
AACTGTTCTTAGATATCCATGCTATTTATGAAGAACAACAGGGATCGAGA
AACAG

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 9 whose identifier is 3789 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YIL059C and has the DNA sequence

TTTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCA
GCTTCCACTAATTTAGATGACTATTTCTCATCTTTGCGTCATCTTCTAACA
CCGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAG
TTGATTTTTATTCCAACAGTAT

The match between the T1 sequence and the C1/C2 sequence is

TTTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCA
 GCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACA
 CCGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAG
 TTGATTTTTATTCCAACA

5

The match between the T2 sequence and the C1/C2 sequence is

TTTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCA
 GCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACA
 CCGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAG
 TTGATTTTTATTCCAACA

10

A C1/C2 short loop on chromosome 12 whose identifier is 5289 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLR301W and has the DNA sequence

15

GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
 TTAGGTATGTAGAATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCAT
 AAAAGGGAATCTGCAATTCTACACAATTCTATAAAATATTATTATCATCGTT
 TTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAG
 CTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACAC
 CGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGT
 TGATTTTTATTCCAACAC

25

The match between the T1 sequence and the C1/C2 sequence is

AGAATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAGGGAA
 TCTGCAATTCTACACAATTCTATAAAATATTATTATCATCGTTTTATATGTTA
 ATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCCACTAA
 TTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTATATGAT

30

AATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTTTTATT
CCAACA

The match between the T2 sequence and the C1/C2 sequence is

5

AGGATTTAGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTATA
AATATTATTATCATCGTTTTATATGTTAATATTCATTGATCCTATTACATTA
TCAATCCTTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCATT
TGCCTCATCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATACT
10 AGTTAGTAGATGATAGTTGATTTTTATTCCAACA

A C1/C2 short loop on chromosome 13 whose identifier is 5753 controls the
expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed
as a RNA single strand that is 3'UTR to the gene YMR044W and has the DNA
sequence

15

TTGAGAAATGGGGGAATGTTGAGATAATTGTTGGGATTCCATTGTTGATA
AAGGCTATAATATTAGGTATACAGAATATACTAGAAGTTCTCCTCAAGGA
TATAGGAATCCTCAAAATGGAATCTATTTCTACATACTAATATTACGAT
20 TATTCCTCATTCCGTTTTATATGTTTCATTATCCTATTACATTATCAATCCT
TGCACTTCAGCTTCCTCTAACTTCGATGACAGCTTCTCATAACTTATGTCA
TCATCTTAACACCGTATATGATAATATATTGATAATATAACTATTAGTTGA
TAGACGATAGTGGATTTTTATTCCAACAT

25

The match between the T1 sequence and the C1/C2 sequence is

AGATAATTGTTGGGATTCCATTGTTGATAAAGGCTATAATATTAGGTATAC
AGAATATACTAGAAGTTCTCCTC

30

The match between the T2 sequence and the C1/C2 sequence is

09060203040506070809101112131415161718192021222324252627282930


TTGTTGGGATTCCATTGTTGATAAAGGCTATAATATTAGGTATACAGAATA
TACTAGAAGTTCTCCTCAAGGAT

5

Two examples of multi-celled geneless connectrons – *C. elegans*

In the first example the existence of the T1-T2 (2342-2344) long loop is controlled by the C1/C2 (24114) short loop. The expression of one C1/C2 (2343) short loop is controlled by the existence of the T1-T2 (2342-2344) long loop.

10




15

20

In the second example the existence of the T1-T2 (29221-29262) long loop is controlled by the C1/C2 (24114) short loop. The expression of one C1/C2 (2343) short loop is controlled by the existence of the T1-T2 (2342-2344) long loop.

25



30

35

A double stranded DNA loop of length 67.059 kilo-bases on chromosome 1 is bounded on the left by a T1 sequence whose identifier is 2342. This T1 control element has the DNA sequence

TGAAAACACTACAGTAATTCTTTAAATGACTACTGTAGC

5 This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 2344. This T2 control element has the DNA sequence

CTACTGTAGCGCTTGTGTCGATTTACGGGCTCGATTT

10 There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15 A C1/C2 short loop on chromosome 1 whose identifier is 2343 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

TCGACACAAGCGCTACAGTAGCTATTTAAAGAATTACTGTAGTTTTTCGCTA
CGAGATATTT

20 The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 5 whose identifier is 24114 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene C13F10.5 and has the DNA sequence

30 GCGAAAACACTACAGTAATTCTTTAAATGACTACTGTAGCGCTTGTGTCGATT
TACGGGCTCGATTTTCG

The match between the T1 sequence and the C1/C2 sequence is

GAAAACTACAGTAATTCTTTAAATGACTACTGTAGC

The match between the T2 sequence and the C1/C2 sequence is

CTACTGTAGCGCTTGTGTCGATTTACGGGCTCGATT

A double stranded DNA loop of length 41.297 kilo-bases on chromosome 5 is bounded on the left by a T1 sequence whose identifier is 29221. This T1 control element has the DNA sequence

TTTAAATTTCCCGCCAAAAATTGACTGAAAATTTGGATTTTCTTTCCAAAA
ATTGACAGAAA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 29262. This T2 control element has the DNA sequence

TGAAAATTTGAATTTCCCGCCAAAAATTAAC

There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 5 whose identifier is 29222 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

AATTTCCCGCCAAAAATTGACTGAAAATTTGGATTTTCTTTCCAAAAATTG
ACAGAAA

This T1-T2 loop also modulates the C1/C2 short loops numbered 29223 to 29260

A C1/C2 short loop on chromosome 5 whose identifier is 29261 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

AAAATTGACTGAAAATTTGAATTTCCAGCCAAAAATTGACTGAAAATTTG
AATT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 1 whose identifier is 4291 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene Y43F8C.5 and has the DNA sequence

AAAATTAAGTAAAATTTGAATTTCCCGCCAAAAATTGACTGAAAATTTG
AATTTCCCGCCAAAAAAATTGACTGAAAATTTGAATTTCCCGCCAAAAA
TTGACTGAAAATTTGAATTTCCCGCCAAAAATTAATTGAAAATTTGAATTT
CCCGCCAAAAATTAATTGAACTTTGAATTTTCAA...ATTTCCCGCCAAAA
ATTAATTGAACTTTGAATTTTCAAATTTCCCGCCAAAAATTGACTGAAAA
TTTGAATTTCCCGCCAAAAATTAATTGAAAATTTGAATTTTGAATTTCC
GCCAAAAATGACTGA

The match between the T1 sequence and the C1/C2 sequence is

TTTAAATTTCCCGCCAAAAATTGACTGAAAATTTG

The match between the T2 sequence and the C1/C2 sequence is

AAAAAAATTGACTGAAAATTTGAATTTCCCGCCAAAAATTGA


0966925.053004

10. One connectron controls many geneless connectrons in single-celled and multi-celled eukaryotes

One C1/C2 short loop can control the existence of many geneless T1-T2 long loops.

Example of a single-celled geneless connectron – *S. cerevisiae*

In this example the existence of the three T1-T2 (1142-1156, 1242-1272 and 7102-7117) long loops is controlled by the C1/C2 (5289) short loop.



A double stranded DNA loop of length 5.337 kilo-bases on chromosome 4 is bounded on the left by a T1 sequence whose identifier is 1142. This T1 control element has the DNA sequence

ATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATATTAGG
TATGTAGATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAG

GGAATCTGCAATTCTACACAATTCTATAAAATATTATTATCATCATTTTATA
 TGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCC
 ACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTAT
 ATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTT
 TTATTCCAACA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1156. This T2 control element has the DNA sequence

TTTTAATAAGGCAATAATATTAGGTATGTAGATATACTAGAAGTTCTCCTC
 CAGGATTTAGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTAT
 AAATATTATTATCATCATTTTATATGTTAATATTCATTGATCCTATTACATT
 ATCAATCCTTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCAT
 TTGCGTCATCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATAC
 TAGTTAGTAGATGATAGTTGATTTTTATTCCAACAAGAA

There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 4 whose identifier is 1143 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

ATTTTGAGATAATTGTTGGGATTCCATTTTTAATAAGGCAATAATATTAGG
 TATGTAGATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAG
 GGAATCTGCAATTCTACACAATTCTATAAAATATTATTATCATCATTTTATA
 TGTTAATATTCATTGATCCTATTACATTATCAAT...CTCTAAGTCTCATTGCC
 TTTGTGCCAAAAAATCTGTTTCTAAATTTCTCTTCATTTGTAGACTTAATTA
 TACTGATCGTTGATCTACTATCAGTAAGTAAGCCTTTAATAATTGGTTTCT
 TGTTAAGTTCTTGCACAAGGTGACTGAGGTTATTCAATAGCGG

This T1-T2 loop also modulates the C1/C2 short loops numbered 1144 to 1154

A C1/C2 short loop on chromosome 4 whose identifier is 1155 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

GAGGAGAACTTCTAGTATATCTACATACCTAATATTATTGCCTTATTA
ATGGAATCCCAACAATTA

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 12 whose identifier is 5289 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLR301W and has the DNA sequence

GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
TTAGGTATGTAGAATATACTAGAAAGTTCTCCTCGAGGATTTAGGAATCCAT
AAAAGGGAATCTGCAATTCTACACAATTCTATAAATATTATTATCATCGTT
TTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAG
CTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACAC
CGTATATGATAATATACTAGTACGTAAATACTAGTTAGTAGATGATAGTT
GATTTTTATTCCAACAC

The match between the T1 sequence and the C1/C2 sequence is

ATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATATTAGG
TATGTAGA

The match between the T2 sequence and the C1/C2 sequence is

TTTTAATAAGGCAATAATATTAGGTATGTAGA

5

A double stranded DNA loop of length 5.251 kilo-bases on chromosome 4 is bounded on the left by a T1 sequence whose identifier is 1243. This T1 control element has the DNA sequence

10

CGTGTTTTATCTCATGTTGTTTCGTTTTGTTATTGAGATATATGTGGGTAATT
AGATAATTGTTGGGATTCCATTGTTGATAAAGGCTATAATATTAGGTATAC
AGAATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAGGGAA
TCTGCAATTCTACACAATTCTATAAAATATTATTATCATCGTTTTATATGTTA
ATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCCACTAA
TTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTATATGAT
AATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTTTTATT
CCAACA

15

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 1272. This T2 control element has the DNA sequence

20

TGAGATATATGTGGGTAATTAGATAATTGTTGGGATTCCATTGTTGATAAA
GGCTATAATATTAGGTATACAGAATATACTAGAAGTTCTCCTCGAGGATTT
AGGAATCCATAAAAGGGAATCTGCAATTCTACACAATTCTATAAAATATTA
TTATCATCGTTTTATATGTTAATATTCATTGATC...TATACTAGTAACGTAA
ATACTAGTTAGTAGATGATAGTTGATTTTTATTCCAACAGTTATAAGGTTG
TTTCATATGTGTTTTATGAA

25

There are no genes controlled by this T1/T2 loop.

30

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 4 whose identifier is 1244 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

5

TTTATCTCATGTTGTTTCGTTTTGTTATTGAGATATATGTGGGTAATTAGATA
ATTGTTGGGATTCCATTGTTGATAAAGGCTATAATATTAGGTATACAGAAT
ATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAGGGAATCTGC
AATTCTACACAATTCTATAAATATTATTATCAT...GTCTCGATGTAGTATAC
GTATAAATTATTACCTGATACTTCATCTCTAAGTCTCATTGCCTTTGTGCCA
AAAAATCTGTTTCTAAATTTCTCTTCATTTGTAGACTTAATTATACTGATCG
TTGATCTACTATCAGTAAGT

10

This T1-T2 loop also modulates the C1/C2 short loops numbered 1245 to 1270

15

A C1/C2 short loop on chromosome 4 whose identifier is 1271 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

20

TGTTGTATCTCAAAATGAGATATGTCAGTATGACAATACGTCATCCTAAAC
GTTCATAAAACACATATGAAACAACCTTATAACTGTTGGAATAAAAATCA
ACTATCATCTACTAACTAGTATTTACGTTACTAGTATATTATCATATACGG
TGTTAGAAGATGACGCAAATGATGAGAAATAGTC...CAACAATGGAATCC
CAACAATTATCTAATTACCCACATATATCTCATGGTAGCGCCTGTGCTTCG
GTTACTTCTAAGGAAGTCCACACAAATCAAGATCCGTTAGACGTTTCAGC
TTCCAAAA

25

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

30

A C1/C2 short loop on chromosome 12 whose identifier is 5289 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed

as a RNA single strand that is 3'UTR to the gene YLR301W and has the DNA sequence

5 GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
TTAGGTATGTAGAATATACTAGAAAGTTCTCCTCGAGGATTTAGGAATCCAT
AAAAGGGAATCTGCAATTCTACACAATTCTATAAATATTATTATCATCGTT
TTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAG
CTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACAC
CGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGT
10 TGATTTTATTCCAACAC

The match between the T1 sequence and the C1/C2 sequence is

15 AGAATATACTAGAAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAGGGAA
TCTGCAATTCTACACAATTCTATAAATATTATTATCATCGTTTTATATGTTA
ATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCCACTAA
TTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTATATGAT
AATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTTTATT
CCAACA

The match between the T2 sequence and the C1/C2 sequence is

25 AGAATATACTAGAAAGTTCTCCTCGAGGATTTAGGAATCCATAAAAGGGAA
TCTGCAATTCTACACAATTCTATAAATATTATTATCATCGTTTTATATGTTA
ATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCCACTAA
TTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTATATGAT
AATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTTTATT
CCAACA

30 -----

A double stranded DNA loop of length 5.296 kilo-bases on chromosome 15 is bounded on the left by a T1 sequence whose identifier is 7102. This T1 control element has the DNA sequence

CATGATTAATATGACCAATCGGCGTGTGTTTTTGAAAAGTGGGTGAATTTT
GAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATATTAGGTATGT
AGAATGTACTAGAAGTTCTCCTCAAGGATTTAGGAATCCATGAAAGGGAA
TCTGCAATTCTACACAATTCTATAAAATATTATTATCATCATTTTTATATGTTA
ATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAGCTTCCACTAA
TTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACACCGTATATGAT
AATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGTTGATTTTTATT
CCAACA

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 7117. This T2 control element has the DNA sequence

TGAAAAGTGGGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAG
GCAATAATATTAGGTATGTAGAATGTACTAGAAGTTCTCCTCAAGGATTT
AGGAATCCATGAAAGGGAATCTGCAATTCTACACAATTCTATAAAATATTA
TTATCATCATTTTTATATGTTAATATTCATTGATCCTATTACATTATCAATCC
TTGCGTTTCAGCTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCA
TCTTCTAACACCGTATATGATAATATACTAGTAACGTAAATACTAGTTAGT
AGATGATAGTTGATTTTTATTCCAACAGTTTTATATACCTCTCTTATTAGT
ATAAGAA

There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

A C1/C2 short loop on chromosome 15 whose identifier is 7103 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

5 AAGAACATTGCTGATGTGATGACAAAACCTCTTCCGATAAAAAACATTTAA
ACTATTAACAAATGGAATTCATTAGATCTATTACATTATGGGTGGTAT
GTTGGAATAAAAAATCAACTATCATCTACTAACTAGTATTTACGTTACTAGT
ATATTATCATATACGGTGTTAGAAGATGACGCAAATGATGAGAAATAGTC
ATCTAAATTAGTGGAAGCTGAAACGCAAGGATTGATAATGTAATAGGATC
10 AATGAATATTAACATATAAAATGATGATAATAATTTATAGAATTGTGT
AGAATTGCAGATTCCCTTTTCATGGATTCTAAATCCTTGAGGAGAACTTCT
AGTA

This T1-T2 loop also modulates the C1/C2 short loops numbered 7104 to 7115

A C1/C2 short loop on chromosome 15 whose identifier is 7116 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

15 CCATTCTGTGGAGGTGGTACTGAAGCAGGTTGAGGAGAGACATGATGATG
20 GTTCTCTGGAACAGCT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

25 A C1/C2 short loop on chromosome 12 whose identifier is 5289 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene YLR301W and has the DNA sequence

30 GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
TTAGGTATGTAGAATATACTAGAAGTTCTCCTCGAGGATTTAGGAATCCAT

AAAAGGGAATCTGCAATTCTACACAATTCTATAAAATATTATTATCATCGTT
 TTATATGTTAATATTCATTGATCCTATTACATTATCAATCCTTGCGTTTCAG
 CTTCCACTAATTTAGATGACTATTTCTCATCATTTGCGTCATCTTCTAACAC
 CGTATATGATAATATACTAGTAACGTAAATACTAGTTAGTAGATGATAGT
 TGATTTTTATTCCAACAC

The match between the T1 sequence and the C1/C2 sequence is


GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
 TTAGGTATGTAGAAT

The match between the T2 sequence and the C1/C2 sequence is

GGTGAATTTTGAGATAATTGTTGGGATTCCATTTTAAATAAGGCAATAATA
 TTAGGTATGTAGAAT

Example of a multi-celled geneless connectron – *C. elegans*

In this example the existence of the three T1-T2 (1142-1156, 14840-15042 and 15365-15627) long loops is controlled by the C1/C2 (16760) short loop.



A C1/C2 short loop on chromosome 1 whose identifier is 3119 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

CCCGCATTTTTGTAGATCAAACCGTAATGGGACGGCCTGGCAACACGTG
ATTTTCCAAAT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 4 whose identifier is 16760 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene T23E1.2 and has the DNA sequence

GGCAAATTGCCGAAATTGAACATTTCCGGCAAATCGGCAAATTGCCGGAA
TTGAACATTTCCGGCAAATCGGCAAATTGCCGGAATTGAACATTTCCGGC
AAATCGGCAAATTGCCGGAATTGA

The match between the T1 sequence and the C1/C2 sequence is

CAAATCGGCAAATTGCCGGAATTGAACATTTCC

The match between the T2 sequence and the C1/C2 sequence is

TTTCCGGCAAATCGGCAAATTGCCGGAATTG

A double stranded DNA loop of length 86.977 kilo-bases on chromosome 3 is bounded on the left by a T1 sequence whose identifier is 14840. This T1 control element has the DNA sequence

AAAAATTTCCGGCAAGTCGGCAATTTTCCGAAAATGAAAATTTCCGGCAA
ATCGGCAAATTGCCGGAATTGAAAATTCCTGGCAAATCAGCAAATTTGCG
GCAAATCGGCAATTTGCCGAAAATGAAAATTTCCGGCAAAT

5

This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 15042. This T2 control element has the DNA sequence

CAAATCGGTAGGTAAATTGGCCAAACTTGAAAATTTCCGGCAAATCGGCA
AATTCCGCGAACTGAACATTTCCGGCAAATCGGCAAATTGCTCGAACT

10

There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15

A C1/C2 short loop on chromosome 3 whose identifier is 14841 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

AAAAATTTCCGGCAAGTCGGCAATTTTCCGAAAATGAAAATTTCCGGCAA
ATCGGCAAATTGCCGGAATTGAAAATTCCTGGCAAATCAGCAAATTTGCG
GCAAATCGGCAATTTGCCGAAAATGAAAATTTCCGGCAAAT

20

25

This T1-T2 loop also modulates the C1/C2 short loops numbered 14842 to 15040

A C1/C2 short loop on chromosome 3 whose identifier is 15041 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

30

CGGCAATTGCCGTTCCGGCAATTTGCCAATTTGCCGGAAATTTCAATTCCG
GCAA

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

5 A C1/C2 short loop on chromosome 4 whose identifier is 16760 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene T23E1.2 and has the DNA sequence

10 GGCAAATTGCCGAAATTGAACATTTCCGGCAAATCGGCAAATTGCCGGAA
TTGAACATTTCCGGCAAATCGGCAAATTGCCGGAATTGAACATTTCCGGC
AAATCGGCAAATTGCCGGAATTGA

The match between the T1 sequence and the C1/C2 sequence is

15 ATTTCCGGCAAATCGGCAAATTGCCGGAATTGAA

The match between the T2 sequence and the C1/C2 sequence is

20 TGAACATTTCCGGCAAATCGGCAAATTGC

25 A double stranded DNA loop of length 98.488 kilo-bases on chromosome 3 is bounded on the left by a T1 sequence whose identifier is 15365. This T1 control element has the DNA sequence

30 AAAATTTCCGGCAAATCGGCAATTTGCCAAAAATTGAAATTTCCGGCAAA
TCGGCAATTTGTCAAAAATGAAAATTTCCGGCAAATCGGCAAATTGCCGA
AAATGAAAATTTCCGGCAAATCGGCAAACCTCCGGAAGTAAAATTTCCG
GCAAATCGGCAATTTGCCATAAATGAACATTTCCGG...GGCGAAAATTAAA
ATTTCCGCCATATCGGCAATTTGCCAAAAAATTAAAATTTCCGGCAAATC

GGCAAATTGCCGGAATTCAAAATTTCCGGCAAACCGGCAAATTGCCGGAA
CTCAAAATTTCCCGGCAAATCAGCAAATTGCCGGAATT

5 This double stranded DNA loop is bounded on the right by a T2 control element whose identifier is 15627. This T2 control element has the DNA sequence

TGGCAAACCGGCAAATTGCCGGAATTGAACATTTCCGGCAAATCGGCAAT
TTGCCGGAATTGAAATTT

10 There are no genes controlled by this T1/T2 loop.

This long T1/T2 double stranded DNA loop modulates the expression of the following C1/C2 short loops

15 A C1/C2 short loop on chromosome 3 whose identifier is 15366 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

TGCCGATTTGCCGGAAATTTTCATTTTCGGCAATTTGCCGATTTGCCGGAA
20 ATTTTCATT

This T1-T2 loop also modulates the C1/C2 short loops numbered 15366 to 15624

25 A C1/C2 short loop on chromosome 3 whose identifier is 15625 controls the expression of the genes of one or more other T1/T2 long loops. This C1/C2 short loop has the DNA sequence

TCAAGCAAATTGTCAAATTCGCGGAACTAAACATTTCCGGCAAATCGGCA
30 AATT

The expression of genes in this T1/T2 long loop is controlled by the following C1/C2 short loops.

A C1/C2 short loop on chromosome 4 whose identifier is 16760 controls the expression of the genes in this T1/T2 long loop. This C1/C2 short loop is expressed as a RNA single strand that is 3'UTR to the gene T23E1.2 and has the DNA sequence

5

GGCAAATTGCCGAAATTGAACATTTCCGGCAAATCGGCAAATTGCCGGAA
TTGAACATTTCCGGCAAATCGGCAAATTGCCGGAATTGAACATTTCCGGC
AAATCGGCAAATTGCCGGAATTGA

10

The match between the T1 sequence and the C1/C2 sequence is

ATTTCCGGCAAATCGGCAAATTGCCGGAATT

The match between the T2 sequence and the C1/C2 sequence is

15

CGGCAAATTGCCGGAATTGAACATTTCCGGCAAATCGGCAA

T00E50" 23E99B50